

2015

# MRI is the most effective initial diagnostic study for osteomyelitis of the foot in diabetic patients

Anthony M. Provenzano B.A.

Wayne State University, aprovenzano123@gmail.com

Follow this and additional works at: <https://digitalcommons.wayne.edu/crp>

## Recommended Citation

Provenzano AM. MRI is the most effective initial diagnostic study for osteomyelitis of the foot in diabetic patients. *Clin. Res. Pract.* 2015;1(1):eP1000. doi: 10.22237/crp/1422568065

This Meta-analysis Review is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Clinical Research in Practice: The Journal of Team Hippocrates by an authorized editor of DigitalCommons@WayneState.

## META-ANALYSIS REVIEW: MRI is the most effective initial diagnostic study for osteomyelitis of the foot in diabetic patients

ANTHONY M. PROVENZANO, B.A., Wayne State University, Detroit, MI, [aprovenz@med.wayne.edu](mailto:aprovenz@med.wayne.edu)

**ABSTRACT** What is the best initial diagnostic study that should be used for a suspected osteomyelitis diagnosis? A critical review and clinical application of: Kapoor A, Page S, LaValley M, Gale DR, Felson DT. Magnetic resonance imaging for diagnosing foot osteomyelitis: a meta-analysis. *Arch Intern Med.* 2007;167(2):125-132. doi:[10.1001/archinte.167.2.125](https://doi.org/10.1001/archinte.167.2.125). **Keywords:** foot osteomyelitis, MRI, magnetic resonance imaging, diabetes, diagnostic methods, Charcot joint

### Clinical Context

During our infectious disease rounds, we had a patient with suspected osteomyelitis due to a painful toe joint with overriding necrotic ulceration; the patient also had diabetes as comorbidity. Our attending decided to order an MRI (magnetic resonance imaging) as the initial study to work up the possible osteomyelitis. This confused me because at other clerkship sites, we had done bone scans as the initial diagnostic work-up for osteomyelitis.

### Clinical Question

What is the best initial diagnostic study that should be used for a suspected osteomyelitis diagnosis?

### Research Article

Kapoor A, Page S, LaValley M, Gale DR, Felson DT. Magnetic resonance imaging for diagnosing foot osteomyelitis: a meta-analysis. *Arch Intern Med.* 2007;167(2):125-132. doi:[10.1001/archinte.167.2.125](https://doi.org/10.1001/archinte.167.2.125).

### Literature Review

The above-identified research article, provided by a mentor, examined the four most widely used diagnostic tests for osteomyelitis. Keywords derived from this meta-analysis led to multiple prospective trials relevant to the topic of osteomyelitis diagnosis in diabetic patients using MRI. All of the studies included<sup>1-38</sup> had small sample sizes (the largest was only 72)<sup>1</sup>, and many were not designed prospectively nor were they randomized. This meta-analysis has been cited in multiple articles referring to osteomyelitis and is often used as the benchmark study for this subject. This makes it the most relevant article related to imaging techniques in the setting of potential osteomyelitis. Within the meta-analysis, the studies were chosen with the following selection criteria: "Studies were en-rolled when information from the usual diagnostic performance 2x2 table... could be extracted about discrete foot and ankle cases, when 80% or more of the patients were 16 years or older, and when at least one site with the disease and one without were identified by the reference standard." There were 2,053 potential studies excluded from the meta-analysis because

ANTHONY M. PROVENZANO, B.A., is a student in the Wayne State University School of Medicine and Senior Student Editor of *CRP*.



they were unable to meet the criteria mentioned above, or because a patient was reused in two studies. One study was also excluded that included a Charcot joint; this is important because in the article, the authors explain that MRI changes seen in osteomyelitis can be confused when a Charcot joint is present. This is an area of potential selection bias, which will be examined in detail later.

## Critical Appraisal

This meta-analysis provides level 2a evidence using the Oxford and National Guidelines Clearinghouse criteria, because it provides data from cohort and randomized controlled trials. One of the advantages of this study is that it provides a cut-off point for the sensitivity of MRI, which is clinically relevant at 90%. It also provides receiver operator curves with relevant cut points and scales showing the area under curve (AUC) of MRI against the AUC of the three other tests. This kind of direct comparison makes it easy to see the global evidence and superiority of MRI graphically. The drawback of this technique is that it cannot determine the strengths and weaknesses of each individual data set, because it works as an average. A likelihood ratio matrix showing the strengths and weaknesses of each individual study could strengthen this meta-analysis. Instead, the study uses diagnostic odds ratios, thus providing good evidence that the MRI is the best imaging technique to diagnose osteomyelitis of the foot and ankle in adults. Even in subsets that included different designs and more diverse patients, the superior performance of MRI remained statistically relevant. However, there are many flaws in this meta-analysis. For example, very few of the studies used in the meta-analysis followed up the MRI, and corroborated the evidence of osteomyelitis, with the gold standard of biopsy; thus, the comparison of MRI against other techniques for diagnosis was not entirely complete. Also, as noted earlier, the exclusion of documentation regarding Charcot foot is a potential confounding variable since this is often confused with osteomyelitis on MRI. It would be good to establish a best initial diagnostic procedure in this subset of patients.

The largest and most relevant study within this meta-analysis is by Ledermann et al.<sup>1</sup>, which uses standard primary signs (the presence of focally decreased marrow signal on T1 weighted images, and increased marrow signal on T2 weighted images), as well as many secondary signs like ulcers or cortical disruption, on MRI to determine a positive result. The positive likelihood ratio of this study is 5.62 and the negative likelihood ratio is 0.112. These numbers mean that when MRI is positive there is a moderate increase in the likelihood of having osteomyelitis, and when MRI is negative there is moderate-strong decrease in the likelihood of disease.<sup>38</sup> These data agree with the claim put forth in the meta-analysis.

## Clinical Application

After reading this study, it makes sense that we ordered an MRI for my patient because it provides the best initial imaging to diagnose osteomyelitis. Before the MRI results were available, the patient was started on empiric antibiotics. Making a correct diagnosis is essential to avoid continuing antibiotics, which have potentially serious side effects if the patient does not have osteomyelitis. For this patient, the MRI was positive and was confirmed by bone biopsy done for culture and identification of the organism.

Three learning points to share:

- 1.) I learned that bone scans are inferior to MRI for diagnosing osteomyelitis because of excess false positive and false negative results.
- 2.) For my future career, I learned that I would not prescribe antibiotics to patients suspected of having osteomyelitis but who are clinically stable until the MRI results are available.
- 3.) I want to share with my peers that different consultants can give conflicting advice and that ultimately, we as the provider need the ability to make judgment calls regarding appropriate treatment and evaluation plans.



## References

- Ledermann HP, Schweitzer ME, Morrison WB. Nonenhancing tissue on MR imaging of pedal infection: characterization of necrotic tissue and associated limitations for diagnosis of osteomyelitis and abscess. *AJR Am J Roentgenol.* 2002 Jan;178(1):215-22. doi:[10.2214/ajr.178.1.1780215](https://doi.org/10.2214/ajr.178.1.1780215)
- Agency for Healthcare Research Quality, U.S. Department of Health and Human Services. *HCUPnet: Health Care Utilization Project (HCUP)*. Available at <http://hcup.ahrq.gov>, Accessed November 9, 2005.
- Jude EB, Selby PL, Mawer EB, Burgess J, Boulton AJM. Inflammatory and bone turnover markers in Charcot arthropathy and osteomyelitis of the feet in diabetic patients. *Diabetologia* 2002;45(suppl 2):A341-A342.
- Eneroth M, Larsson J, Apelqvist J. Deep foot infections in patients with diabetes and foot ulcer: an entity with different characteristics, treatments, and prognosis. *J Diabetes Complications.* 1999 Sep-Dec;13(5-6):254-63. doi:[10.1016/S1056-8727\(99\)00065-3](https://doi.org/10.1016/S1056-8727(99)00065-3)
- Black ER, Bordley DR, Tape TG, Panzer RJ. *Diagnostic Strategies for Common Medical Problems.* 2nd ed. East Peoria, Ill: Versa Press; 1999.
- Jeffcoate WJ, Lipsky BA. Controversies in diagnosing and managing osteomyelitis of the foot in diabetes. *Clin Infect Dis.* 2004 Aug 1;39 Suppl 2:S115-22. doi:[10.1086/383272](https://doi.org/10.1086/383272)
- Becker W. Imaging osteomyelitis and the diabetic foot. *Q J Nucl Med.* 1999 Mar;43(1):9-20.
- Karchevsky M, Schweitzer ME, Morrison WB, Parellada JA. MRI findings of septic arthritis and associated osteomyelitis in adults. *AJR Am J Roentgenol.* 2004 Jan;182(1):119-22. doi:[10.2214/ajr.182.1.1820119](https://doi.org/10.2214/ajr.182.1.1820119)
- Eckman MH, Greenfield S, Mackey WC, et al. Foot infections in diabetic patients: decision and cost-effectiveness analyses. *JAMA.* 1995 Mar 1;273(9):712-20. doi:[10.1001/jama.1995.03520330042035](https://doi.org/10.1001/jama.1995.03520330042035)
- Matowe L, Gilbert FJ. How to synthesize evidence for imaging guidelines. *Clin Radiol.* 2004 Jan;59(1):63-8. doi:[10.1016/j.crad.2003.09.002](https://doi.org/10.1016/j.crad.2003.09.002)
- Termaat MF, Raijmakers PGHM, Scholten HJ, Bakker FC, Patka P, Haarman HJTM. The accuracy of diagnostic imaging for the assessment of chronic osteomyelitis: a systematic review and meta-analysis. *J Bone Joint Surg Am.* 2005 Nov;87(11):2464-71. doi:[10.2106/JBJS.D.02691](https://doi.org/10.2106/JBJS.D.02691)
- Bossuyt PM, Reitsma JB, Bruns DE, et al. The STARD statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Ann Intern Med.* 2003 Jan 7;138(1):W1-12. doi:[10.7326/0003-4819-138-1-200301070-00012-w1](https://doi.org/10.7326/0003-4819-138-1-200301070-00012-w1)
- Littenberg B, Moses LE. Estimating diagnostic accuracy from multiple conflicting reports: a new meta-analytic method. *Med Decis Making.* 1993 Oct-Dec;13(4):313-21. doi:[10.1177/0272989X9301300408](https://doi.org/10.1177/0272989X9301300408)
- Moses LE, Shapiro D, Littenberg B. Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. *Stat Med.* 1993 Jul 30;12(14):1293-316. doi:[10.1002/sim.4780121403](https://doi.org/10.1002/sim.4780121403)
- Craig JG, Amin MB, Wu K, et al. Osteomyelitis of the diabetic foot: MR imaging—pathologic correlation. *Radiology.* 1997 Jun;203(3):849-55. doi:[10.1148/radiology.203.3.9169715](https://doi.org/10.1148/radiology.203.3.9169715)
- Croll SD, Nicholas GG, Osborne MA, Wasser TE, Jones S. Role of magnetic resonance imaging in the diagnosis of osteomyelitis in diabetic foot infections. *J Vasc Surg.* 1996 Aug;24(2):266-70. doi:[10.1016/S0741-5214\(96\)70102-7](https://doi.org/10.1016/S0741-5214(96)70102-7)
- Enderle MD, Coerper S, Schweizer HP, et al. Correlation of imaging techniques to histopathology in patients with diabetic foot syndrome and clinical suspicion of chronic osteomyelitis: the role of high-resolution ultrasound. *Diabetes Care.* 1999 Feb;22(2):294-9. doi:[10.2337/diacare.22.2.294](https://doi.org/10.2337/diacare.22.2.294)
- Ertugrul MB, Baktiroglu S, Salman S, et al. The diagnosis of osteomyelitis of the foot in diabetes: microbiological examination vs. magnetic resonance imaging and labelled leucocyte scanning. *Diabet Med.* 2006 Jun;23(6):649-53. doi:[10.1111/j.1464-5491.2006.01887.x](https://doi.org/10.1111/j.1464-5491.2006.01887.x)
- Horowitz JD, Durham JR, Nease DB, Lukens ML, Wright JG, Smead WL. Prospective evaluation of magnetic resonance imaging in the management of acute diabetic foot infections. *Ann Vasc Surg.* 1993 Jan;7(1):44-50. doi:[10.1007/BF02042659](https://doi.org/10.1007/BF02042659)
- Kearney T, Pointin K, Cunningham D, Gedroyc W, Robinson S, Elkeles RS. The detection of pedal osteomyelitis in diabetic patients. *Pract Diabetes Int.* 1999; 16(4):98-100. doi:[10.1002/pdi.1960160404](https://doi.org/10.1002/pdi.1960160404)
- Levine SE, Neagle CE, Esterhai JL, Wright DG, Dalinka MK. Magnetic resonance imaging for the diagnosis of osteomyelitis in the diabetic patient with a foot ulcer. *Foot Ankle Int.* 1994 Mar;15(3):151-6.
- Lipman BT, Collier BD, Carrera GF, et al. Detection of osteomyelitis in the neuropathic foot: nuclear medicine, MRI and conventional radiography. *Clin Nucl Med.* 1998 Feb;23(2):77-82.
- Maas M, Slim EJ, Heeksma AF, et al. MR imaging of neuropathic feet in leprosy patients with suspected osteomyelitis. *Int J Lepr Other Mycobact Dis.* 2002 Jun;70(2):97-103.



24. Morrison WB, Schweitzer ME, Batte WG, Radack DP, Russel KM. Osteomyelitis of the foot: relative importance of primary and secondary MR imaging signs. *Radiology.* 1998 Jun;207(3):625-32. doi:[10.1148/radiology.207.3.9609883](https://doi.org/10.1148/radiology.207.3.9609883)
25. Nigro ND, Bartynski WS, Grossman SJ, Kruljac S. Clinical impact of magnetic resonance imaging in foot osteomyelitis. *J Am Podiatr Med Assoc.* 1992 Dec;82(12):603-15. doi:[10.7547/87507315-82-12-603](https://doi.org/10.7547/87507315-82-12-603)
26. Remedios D, Valabhji J, Oelbaum R, Sharp P, Mitchell R. 99mTc-nanocolloid scintigraphy for assessing osteomyelitis in diabetic neuropathic feet. *Clin Radiol.* 1998 Feb;53(2):120-5.
27. Seabold JE, Flickinger FW, Kao SC, et al. Indium-111-leukocyte/technetium-99m-MDP bone and magnetic resonance imaging: difficulty of diagnosing osteomyelitis in patients with neuropathic osteoarthropathy. *J Nucl Med.* 1990 May;31(5):549-56.
28. Vesco L, Boulahdour H, Hamissa S, et al. The value of combined radionuclide and magnetic resonance imaging in the diagnosis and conservative management of minimal or localized osteomyelitis of the foot in diabetic patients. *Metabolism.* 1999 Jul;48(7):922-7. doi:[10.1016/S0026-0495\(99\)90230-5](https://doi.org/10.1016/S0026-0495(99)90230-5)
29. Weinstein D, Wang A, Chambers R, Stewart CA, Motz HA. Evaluation of magnetic resonance imaging in the diagnosis of osteomyelitis in diabetic foot infections. *Foot Ankle.* 1993 Jan;14(1):18-22.
30. Yuh WT, Corson JD, Baraniewski HM, et al. Osteomyelitis of the foot in diabetic patients: evaluation with plain film, 99mTc-MDP bone scintigraphy, and MR imaging. *AJR Am J Roentgenol.* 1989 Apr;152(4):795-800.
31. Ahmadi ME, Morrison WB, Carrino JA, Schweitzer ME, Raikin SM, Ledermann HP. Neuropathic arthropathy of the foot with and without superimposed osteomyelitis: MR imaging characteristics. *Radiology.* 2006 Feb;238(2):622-31. doi:[10.1148/radiol.2382041393](https://doi.org/10.1148/radiol.2382041393)
32. Mushlin AI, Littenberg B. Diagnosing pedal osteomyelitis: testing choices and their consequences. *J Gen Intern Med.* 1994 Jan;9(1):1-7.
33. Newman LG, Waller J, Palestro CJ, et al. Unsuspected osteomyelitis in diabetic foot ulcers: diagnosis and monitoring by leukocyte scanning with indium In 111 oxyquinoline. *JAMA.* 1991 Sep 4;266(9):1246-51. doi:[10.1001/jama.1991.03470090080036](https://doi.org/10.1001/jama.1991.03470090080036)
34. Lipsky BA, Berendt AR, Deery HG, et al. Diagnosis and treatment of diabetic foot infections. *Clin Infect Dis.* 2004 Oct 1;39(7):885-910. doi:[10.1086/424846](https://doi.org/10.1086/424846)
35. Dinnes J, Deeks J, Kirby J, Roderick P. A methodological review of how heterogeneity has been examined in systematic reviews of diagnostic test accuracy. *Health Technol Assess.* 2005 Mar;9(12):1-113, iii. doi:[10.3310/hta9120](https://doi.org/10.3310/hta9120)
36. Centers for Medicare & Medicaid Services. CMS-1501-FC: Changes to the Hospital Outpatient PPS for Calendar Year 2006: Addendum B. *CMS.gov.* July 2006; Available at <http://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/HospitalOutpatientPPS/Hospital-Outpatient-Regulations-and-Notices-Items/CMS052176.html>. Accessed August 24 2006.
37. Centers for Medicare & Medicaid Services. RVU06A: Physician fee schedule relative value files. *CMS.gov.* July 2006; Available at <http://www.cms.hhs.gov/PhysicianFeeSched/PFSRVF/list.asp>. Accessed August 24 2006.
38. Lalkhen AG, McCluskey A. Clinical tests: sensitivity and specificity. *Contin Educ Anaesth Crit Care Pain.* 2008;8(6):221-223. doi:[10.1093/bjaceaccp/mkn041](https://doi.org/10.1093/bjaceaccp/mkn041)

