

4-22-2015

Role of IL-16 in CD4(+) T cell-mediated regulation of relapsing multiple sclerosis

Dusanka S. Skundric

Wayne State University, skundric@wayne.edu

William W. Cruikshank

Boston University

Jelena Drulovic

University of Belgrade

Recommended Citation

Skundric DS, Cruikshank WW, Drulovic J. Role of IL-16 in CD4(+) T cell-mediated regulation of relapsing multiple sclerosis. *J Neuroinflammation*. 2015, Apr 22;12(1):78. doi: 10.1186/s12974-015-0292-x
Available at: http://digitalcommons.wayne.edu/med_neurology/2

This Article is brought to you for free and open access by the Department of Neurology at DigitalCommons@WayneState. It has been accepted for inclusion in Neurology Faculty Publications by an authorized administrator of DigitalCommons@WayneState.

NOTICE IN COMPLIANCE WITH PUBLISHER POLICY: Copyright © 2015 Skundric et al.; licensee BioMed Central. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated. <http://dx.doi.org/10.1186/s12974-015-0292-x>

COMMENTARY

Open Access

Role of IL-16 in CD4⁺ T cell-mediated regulation of relapsing multiple sclerosis

Dusanka S Skundric^{1*}, William W Cruikshank² and Jelena Drulovic³

Abstract

In an important article published in *Nature Medicine*, Liu and colleagues described a novel CD4⁺ FoxA1⁺ regulatory T (Treg) cell population as distinct regulators of relapsing-remitting multiple sclerosis (RRMS) and experimental autoimmune encephalomyelitis (EAE). CD4⁺ FoxA1⁺ Treg cells appear as key regulators of responsiveness to therapy with interferon beta (IFN-β) in RRMS patients. Data indicate that CD4⁺FoxA1⁺ FOXP3⁻ Treg cells develop within the central nervous system (CNS), and a potential of cerebellar granule neurons (CGN) in generation of CD4⁺FoxA1⁺PD-L1^{hi}FOXP3⁻ Treg cells from encephalitogenic CD4⁺ T cells.

A CD4 co-receptor specific ligand, IL-16, governs trafficking and biological properties of CD4⁺ T cells irrespective of their activation state. Functions of IL-16, relevant to Treg cells, include expansion of CD4⁺CD25⁺ T cells in long-term cultures with IL-2, *de novo* induction of FOXP-3 and migration of FOXP-3⁺ T cells. IL-16 is highly conserved across species including human and mouse. CGN and neurons in hippocampus contain neuronal-IL-16 (NIL-16), splice variant of immune IL-16, and express CD4 molecule. In a CD4-dependent manner, IL-16 supports cultured CGN survival.

Concomitant studies of RRMS lesions and corresponding MOG₃₅₋₅₅-induced relapsing EAE in (B6 × SJL)F1 (H-2^{b/s}) mice discovered similar roles of IL-16 in regulation of relapsing disease. In RRMS and EAE relapse, peak levels of IL-16 and active caspase-3 correlated with CD4⁺ T cell infiltration and levels of T-bet, Stat-1(Tyr⁷⁰¹), and phosphorylated neurofilaments of axonal cytoskeleton [NF (M + H) P], suggesting a role of locally produced IL-16 in regulation of CD4⁺ Th1 inflammation and axonal damage, respectively. IL-16 was abundantly present in CD4⁺ T cells, followed by CD20⁺ B, CD8⁺ T, CD83⁺ dendritic cells, and Mac-1⁺ microglia. Apart from lesions, bioactive IL-16 was located in normal-appearing white matter (NAWM) and normal-appearing grey matter (NAGM) in RRMS brain and spinal cord.

A cytokine IL-16 emerges as an important regulator of relapsing MS and EAE. Better understanding of immune cell-neuron interactions mediated by IL-16 will foster development of more specific CD4⁺ T cell subset-targeted therapies to prevent or ameliorate progression of neuroinflammation and axonal and neuronal damage. Translational studies necessitate corresponding EAE models.

Keywords: IL-16, Multiple sclerosis, Cerebellum, CD4⁺ T cells, IFN-β, FoxA1, MOG, EAE, Translational, Therapy

Background

Discovery of a novel lineage of regulatory T (Treg) cells, which are CD4⁺FoxA1⁺, as major regulators of responsiveness to IFN-β therapy in patients with relapsing-remitting multiple sclerosis (RRMS), is important for further optimization of this first in class disease-modifying therapy (DMT). Concurrent clinical and experimental evidence support the notion that regulation of MS progression engages distinct regulatory pathway

carried by CD4⁺FoxA1⁺ FOXP3⁻ Treg cells, which develop within the central nervous system (CNS) [1]. To gain better understanding of complex cellular and molecular interactions important for disease progression, some fundamental mechanisms underlying MS immune pathogenesis, reinforced by this recent finding deserve additional attention and reappraisal.

Primarily, a potential of locally inflamed CNS micro-environment to regulate plasticity of CD4⁺ T cell responses, by switching their functional phenotypes from encephalitogenic to suppressive, regulatory, FoxA1 mediated, appears vital for progression of the disease [1]. Locally elaborated cytokines are critical regulators of

* Correspondence: Skundric@wayne.edu

¹Department of Immunology and Microbiology, Wayne State University, 540 East Canfield Av., Detroit, MI, USA

Full list of author information is available at the end of the article

types of inflammatory cell interactions and mediators of tissue damage. In progressive inflammatory disease, where inflammatory mediators are reoccurring, immune cell communications are likely diverse engaging in distinctive types of relationships throughout RRMS and its conversion into secondary progressive (SP) disease.

Additionally, effects of IFN- β therapy regulated through CD4⁺ FoxA1⁺ Treg are shared mechanisms between RRMS and EAE [1]. Such approach further emphasizes importance of an appropriate EAE model when studying immune regulation of MS. In order to identify ongoing modifications in intertwined shared regulatory mechanisms, it is critical for an EAE model to resemble the RRMS clinical course, histopathology and relevant immune mechanisms [2,3]. Immune regulation of relapse in response to myelin oligodendrocyte glycoprotein peptide 35 to 55 (MOG₃₅₋₅₅) is poorly studied due to a relative deficiency of relapsing EAE in mice [4]. MOG is conserved across species. An immunodominant epitope of MOG₃₅₋₅₅ is associated with autoimmune responses by CD4⁺ Th1 and Th17 cells and/or autoantibodies in RRMS and aquaporin-4⁻ (AQP4⁻) relapsing neuromyelitis optica (NMO), respectively [5].

Finally, data from co-culture experiments prompt us to question the nature of interactions between cerebellar granule neurons and encephalitogenic CD4⁺ T cells, in generation of CD4⁺FoxA1⁺PD-L1^{hi}FOXP3⁻ Tregs [1]. Demyelinating lesions in cerebellar cortex are prominent in progressive MS [6]. Clinical signs of cerebellar dysfunction including nystagmus and ataxic gait accompany MS [7]. In homeostasis, hippocampal and cerebellar granule neurons produce neuronal-IL-16 (NIL-16), a splice variant of immune IL-16, which is produced by lymphocytes, macrophages, DC, mast cells, and microglia [8-10].

Commentary

Data from RRMS tissue analysis and relapsing EAE are concurrent in identifying a CD4⁺ T cell-specific chemotactic factor, IL-16, as a key regulator of progressive CNS inflammation mediated by CD4⁺ Th1 cells.

In RRMS acute brain lesions, IL-16 was found in CD3⁺ and CD4⁺ T cells, followed by CD20⁺ B, CD8⁺ T and CD83⁺ dendritic cells and Mac-1⁺ microglia. Often, IL-16 located to cell cytoplasm, less frequently to cell nucleus, or polarized on the membrane connecting adjoining cells. The majority of CD4⁺ IL-16⁺ cells were T-bet⁺ and active caspase-3⁺. Increased levels of bioactive IL-16 correlated with levels of active caspase-3, T-bet, Stat-1 (Tyr⁷⁰¹), and phosphorylated medium and heavy neurofilaments of axonal cytoskeleton, [NF (M + H) P], which suggested a role for IL-16 in regulation of CD4⁺ Th1 inflammation and damage of axonal cytoskeleton, respectively. IL-16 induces migration of CD4⁺Tbet⁺ (Th1) and CD4⁺FOXP3⁺ (Treg) cells and FOXP3 *de novo* expression [11,12]. In the

MS brain, highest levels of IL-16 were detected in chronic lesions, followed by subacute and acute. Conversely, in MS spinal cord, acute lesions contained highest levels of IL-16. Bioactive IL-16 was also detected in normal-appearing white matter (NAWM) and normal-appearing grey matter (NAGM) in the brain and NAWM in the spinal cord. More robust increase of IL-16 was noted in the spinal cord compared to brain NAWM. A sharp increase of pro-IL-16 was found in NAGM in RRMS brain [2]. Data propose regional differences in IL-16 regulation in lesions between the brain and spinal cord and a role of bioactive and pro-IL-16 in pathology of NAWM and NAGM in RRMS.

Corresponding to our findings of elevated IL-16 levels *in situ*, raised levels of IL-16 were detected in serum of MS patients. Following therapy with IFN- β 1a, IL-16 gene expression in peripheral blood mononuclear cells (PBMC) and its serum levels were down regulated, suggesting IL-16 as a potential biomarker for MS [13,14]. Myelin oligodendrocyte glycoprotein (MOG)-specific CD4⁺Th17 exhibited more pronounced IL-16 production compared to CD4⁺Th1 cells, implicating a role of IL-16 in the regulation of Th17 responses [14].

Concurrent with RRMS findings, in relapsing-remitting MOG₃₅₋₅₅-induced EAE in (B6 \times SJL) F1 (H-2^{b/s}) mice, intra-CNS production of IL-16 correlates with extensive CD4⁺ T cell infiltration, accompanied by phosphorylation of axonal cytoskeleton, which all peak during relapses and in chronic progressive disease. IL-16 and CD4 were coprecipitated from CNS of relapsing H-2^{b/s} mice. Data indicate that active caspase-3-mediated release of bioactive IL-16 from infiltrating CD4⁺ T cells supports progression of local inflammation by specifically enhancing trafficking of additional CD4⁺ T cells [2,3,15].

As opposed to low/non-relapsing parental, B6 (H-2^b), in H-2^{b/s} mice, immunization with MOG₃₅₋₅₅ induces relapsing-remitting EAE with extensive CD4⁺ T cell infiltration, followed by Mac-3⁺ macrophages, B220⁺ B, and CD8⁺ T cells. Induced autoimmune responses to MOG₃₅₋₅₅ in H-2^{b/s} but not in H-2^b mice result in myelin-associated glycoprotein (MAG) predominated pattern of demyelination, which is suggestive of oligodendrocyte dysfunction and/or damage [15,16]. A similar pattern of demyelination was observed in the MS lesion subtype III, where profound loss of MAG serves as an indicator of a distal oligodendroglial pathology [17]. MAG-predominated demyelination, a decrease of NF160 axonal neurofilament, and a sharp elevation of PARPp85 suggested axonal dysfunction and irreversible apoptosis, respectively, and correlated with severe relapsing disease in H-2^{b/s} mice [16]. Relapsing-remitting EAE in H-2^{b/s} mice may provide valuable insights into mechanisms of MAG depletion and oligodendroglial pathology initiated by autoimmune responses to MOG₃₅₋₅₅. Therapy with IL-16-specific antibody diminished paralysis

and CD4⁺ T infiltration and abrogated demyelination and axonal damage in relapsing EAE H-2^{b/s} mice [18].

Studies from RRMS and corresponding MOG_{35–55}-induced relapsing EAE in H-2^{b/s} mice suggest a role of IL-16 in neuron-immune cell communications in lesions of gray matter, including cerebellar and hippocampal, and in adjacent NAGM and NAWM (Skundric, unpublished). Cortical and subcortical, including hippocampal, grey matter lesions underlie cognitive deficits observed in MS [19].

Beneficial effects of IFN- β therapy include potential lowering of probability of disease progression from RRMS to SP and development of Expanded Disability Status Scale (EDSS) score [20].

Uncovering those IL-16-specific mechanisms is critical for the development of new CD4⁺ T cell subset-targeted therapies for MS and other chronic and/or progressive inflammatory and demyelinating diseases.

Conclusion

Data from FoxA1 co-culture experiments argue for the importance of bidirectional neuron-immune cell communications and underscore necessity for better understanding of underlying mechanisms, in particular IL-16 mediated. Future studies in this direction are required for the development of specifically targeted therapies aimed at preventing or ameliorating neuronal damage.

Delineating differences in mechanisms of immune regulation between the spinal cord and brain is critical for therapy of NMO and NMO-spectrum disorders (NMOSD) with predominate optic nerve and spinal cord demyelination.

Delineating molecular mechanisms enhancing migration of encephalitogenic CD4⁺ cells (Th1 and Th17), and Treg cells into the CNS throughout the disease, is essential in the development of new therapy specific for CD4⁺ T cell subsets for MS and similar inflammatory and demyelinating diseases.

Abbreviations

[NF(M + H)]P: phosphorylated medium and heavy chains of neurofilament; AQP4: aquaporin-4; DMT: disease-modifying therapy; EAE: experimental autoimmune encephalomyelitis; IFN- β : interferon beta; IL-16: interleukin 16; MAG: myelin-associated glycoprotein; MOG: myelin oligodendrocyte glycoprotein; NAGM: normal-appearing grey matter; NAWM: normal-appearing white matter; NIL-16: neuronal IL-16; NMO: neuromyelitis optica; NMOSD: NMO-spectrum disorders; PBMC: peripheral blood mononuclear cells; RRMS: relapsing-remitting multiple sclerosis; Stat-1: signal transducer and activator of transcription-1.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DSS conceived idea and wrote the commentary. JD contributed with critical comments in regard to IFN- β therapy. WWC contributed with critical comments related to IL-16. All authors read and approved the final manuscript.

Acknowledgements

Data discussed in this commentary was partly supported by the Pilot projects from the National Multiple Sclerosis Society (NMSS) PP0701 and PP0958 and by the Parker Weber Endowment for the Department of Neurology and Detroit Medical Center to DSS.

Author details

¹Department of Immunology and Microbiology, Wayne State University, 540 East Canfield Av., Detroit, MI, USA. ²Pulmonary Center Boston University, 715 Albany Street, Boston, MA, USA. ³Clinic of Neurology, Clinical Center of Serbia, Faculty of Medicine, University of Belgrade, Dr Subotica 6, Belgrade, Serbia.

Received: 30 December 2014 Accepted: 31 March 2015

Published online: 22 April 2015

References

- Liu Y, Carlsson R, Comabella M, Wang J, Kosicki M, Carrion B, et al. FoxA1 directs the lineage and immunosuppressive properties of a novel regulatory T cell population in EAE and MS. *Nat Med*. 2014;20(3):272–82. doi:10.1038/nm.3485. Epub 2014 Feb 16.
- Skundric DS, Cai J, Cruikshank WW, Gveric D. Production of IL-16 correlates with CD4⁺ Th1 inflammation and phosphorylation of axonal cytoskeleton in multiple sclerosis (MS) Lesions. *J Neuroinflammation*. 2006;3(1):13. <http://www.jneuroinflammation.com/content/3/1/13>.
- Skundric DS, Zhou W, Cruikshank WW, Dai R. Increased levels of bioactive IL-16 correlate with disease activity during relapsing experimental autoimmune encephalomyelitis (EAE). *J Autoimmun*. 2005;25(3):206–14.
- Skundric DS. Experimental models of relapsing-remitting multiple sclerosis: current concepts and perspective. *Curr Neurovasc Res*. 2005;2(4):349–62. Review.
- Reindl M, Pauli F, Rostasy K, Berger T. The spectrum of MOG antibody-associated demyelinating diseases. *Nat Rev Neurol*. 2013;9(8):455–61. doi: 10.1038/nrneuro.2013.118.
- Kutzelnigg A, Faber-Rod JC, Bauer J, Lucchinetti CF, Sorensen PS, Laursen H, et al. Widespread demyelination in the cerebellar cortex in multiple sclerosis. *Brain Pathol*. 2007;17:38–44. doi:10.1111/j.1750-3639.2006.00041.
- Damasceno A, Damasceno BP, Cendes F. The clinical impact of cerebellar grey matter pathology in multiple sclerosis. *Plos One*. 2014;9(5):e96193.
- Cruikshank WW, Kornfeld H, Center DM. Interleukin-16. *J Leukoc Biol*. 2000;67(6):757–66.
- Kurchner C, Yuzaki M. Neuronal interleukin-16 (NIL-16): a dual function PDZ domain protein. *Neurosci*. 1999;19:7770–80.
- Croq F, Vizioli J, Tuzova M, Tahtouh M, Sautiere PE, Van Camp C, et al. A homologous form of human interleukin 16 is implicated in microglia recruitment following nervous system injury in leech *Hirudo medicinalis*. *Glia*. 2010;58(14):1649–62. doi:10.1002/glia.21036.
- Lynch EA, Heijens CA, Horst NF, Center DM, Cruikshank WW. Cutting edge: IL-16/CD4 preferentially induces Th1 cell migration: requirement of CCR5. *J Immunol*. 2003;171(10):4965–8.
- McFadden C, Morgan R, Rahangdale S, Green D, Yamasaki H, Center D, et al. Preferential migration of T regulatory cells induced by IL-16. *J Immunol*. 2007;179(10):643–5.
- Annibaldi V, Di Giovanni S, Cannoni S, Giugni E, Bompreszi R, Mattei C, et al. Gene expression profiles reveal homeostatic dynamics during interferon-beta therapy in multiple sclerosis. *Autoimmunity*. 2007;40(1):16–22.
- Nischwitz S, Faber H, Sämman PG, Domingues HS, Krishnamoorthy G, Knop M, et al. Interferon β -1a reduces increased interleukin-16 levels in multiple sclerosis patients. *Acta Neurol Scand*. 2014;130(1):46–52. doi:10.1111/ane.12215.
- Skundric DS, Zakarian V, Dai R, Lisak R, James J. Distinct immune regulation of the response to H-2^b restricted epitope of MOG causes relapsing-remitting EAE in H-2^{b/s} mice. *J Neuroimmunol*. 2003;136:34–45.
- Skundric DS, Zakarian VL, Dai R, Zhou W. Autoimmune-induced preferential depletion of myelin-associated glycoprotein (MAG) is genetically regulated in relapsing EAE (B6 x SJL) F1 mice. *Mol Neurodegener*. 2008;3(1):7. <http://www.molecularneurodegeneration.com/content/3/1/7>.
- Lassmann H, Bruck W, Lucchinetti C. Heterogeneity of multiple sclerosis pathogenesis: implications for diagnosis and therapy. *Trends Mol Med*. 2001;7:115–21.

18. Skundric DS, Dai R, Zakarian VL, Bessert D, Skoff RP, Cruikshank WW, et al. Anti-IL-16 therapy reduces CD4+ T-cell infiltration and improves paralysis and histopathology of relapsing EAE. *J Neurosci Res.* 2005;79(5):680–93.
19. Messina S, Patti F. Gray matters in multiple sclerosis: cognitive impairment and structural MRI. *Mult Scler Int.* 2014;2014:609694.
20. Drulovic J, Kostic J, Mesaros S, Dujmovic Basuroski I, Stojsavljevic N, Kistic-Tepavcevic D, et al. Interferon-beta and disability progression in relapsing-remitting multiple sclerosis. *Clin Neurol Neurosurg.* 2013;115 Suppl 1:S65–9. doi:10.1016/j.clineuro.2013.09.024.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

