

3-30-2021

## I Lost My Taste And Smell... Do I Have COVID-19?

Lauren Kasmikha

Wayne State University School Of Medicine, ft8931@wayne.edu

Dana Labuda

Wayne State University School Of Medicine, he7783@wayne.edu

Ruchi Mangal

Wayne State University School Of Medicine, fz3969@wayne.edu

Brandon Askar

Wayne State University School Of Medicine, fv7083@wayne.edu

Samuel Cain

Wayne State University School Of Medicine, hf0705@wayne.edu

*See next page for additional authors*

## Creative Commons License:



This work is licensed under a [No Rights Reserved](#) license.

---

### Recommended Citation

Kasmikha, Lauren; Labuda, Dana; Mangal, Ruchi; Askar, Brandon; Cain, Samuel; and Rubino, Cameron, "I Lost My Taste And Smell... Do I Have COVID-19?" (2021). *COVID-19 Medical Myth Infographics*. 22. <https://digitalcommons.wayne.edu/covidinfographics/22>

This Infographic is brought to you for free and open access by the Open Source Medicine at DigitalCommons@WayneState. It has been accepted for inclusion in COVID-19 Medical Myth Infographics by an authorized administrator of DigitalCommons@WayneState.

---

## I Lost My Taste And Smell... Do I Have COVID-19?

Follow this and additional works at: <https://digitalcommons.wayne.edu/covidinfographics>



Part of the [Curriculum and Instruction Commons](#), [Medical Education Commons](#), and the [Public Health Commons](#)

### Authors

Lauren Kasmikha, Dana Labuda, Ruchi Mangal, Brandon Askar, Samuel Cain, and Cameron Rubino

# I LOST MY TASTE AND SMELL...

## Do I have COVID-19?


50% of COVID-positive patients had either a loss of taste, smell, or both.

Loss or changes in taste or smell usually last 2-4 weeks, but can be longer

Patients with COVID-19 were 4x more likely to have changes in their taste and smell compared to those with other respiratory viruses.

In one study, 62% of patients over the age of 60 had changes in smell.

COVID-positive patients are 48x more likely to have changes in smell compared to healthy patients.



\*Changes in smell can be complete or partial loss of smell, or a persistent bad/atypical odor.  
\*A loss of taste or smell could be a strong indicator of risk for COVID-19 infection.