

3-30-2021

Ibuprofen Use In COVID-19

Husain Al-Saffar

Wayne State University School Of Medicine, fz3421@wayne.edu

Kriti Babu

Wayne State University School Of Medicine, fw6416@wayne.edu

Kelsey Gockman

Wayne State University School Of Medicine, hf9942@wayne.edu

John Kulka

Wayne State University School Of Medicine, gm7925@wayne.edu

Lauren Wylie

Wayne State University School Of Medicine, ga4762@wayne.edu

Creative Commons License:



This work is licensed under a [No Rights Reserved](#) license.

Recommended Citation

Al-Saffar, Husain; Babu, Kriti; Gockman, Kelsey; Kulka, John; and Wylie, Lauren, "Ibuprofen Use In COVID-19" (2021). *COVID-19 Medical Myth Infographics*. 23.
<https://digitalcommons.wayne.edu/covidinfographics/23>

This Infographic is brought to you for free and open access by the Open Source Medicine at DigitalCommons@WayneState. It has been accepted for inclusion in COVID-19 Medical Myth Infographics by an authorized administrator of DigitalCommons@WayneState.

Ibuprofen Use In COVID-19

Follow this and additional works at: <https://digitalcommons.wayne.edu/covidinfographics>



Part of the [Curriculum and Instruction Commons](#), [Medical Education Commons](#), and the [Public Health Commons](#)

IBUPROFEN USE IN COVID-19

MYTH

Taking Ibuprofen to treat COVID-19 will increase the severity of infection.

TRUTH

Ibuprofen can be used to safely manage symptoms of COVID-19.



Multiple studies compared the use of Ibuprofen (Advil/Motrin) and Acetaminophen (Tylenol) in patients with COVID-19, and saw *NO differences in mortality rates OR the need for respiratory support.*

WHAT YOU SHOULD DO



Follow your doctor's recommendations



Don't change medication use without contacting your physician



Stay home if you are sick or have been in contact with someone who is sick



Manage symptoms with over the counter medications, such as Tylenol or Ibuprofen



Point your phone camera to the code on the left to view current CDC guidelines regarding COVID-19.