

4-1-2016

Paternal Genetic Structure In Contemporary Mennonite Communities From The American Midwest

Kristine G. Beaty

Laboratories of Biological Anthropology, University of Kansas, Lawrence, KS, kgbeaty@gmail.com

M. J. Mosher

Department of Anthropology, Western Washington University, Bellingham, WA, m.j.mosher@wwu.edu

Michael H. Crawford

Laboratories of Biological Anthropology, University of Kansas, Lawrence, KS, crawford@ku.edu

Phillip Melton

Curtin/UWA Centre for Genetic Origins of Health and Disease, Faculty of Health Sciences, Curtin University, and Faculty of Medicine Dentistry and Health Sciences, University of Western Australia, Perth, Australia, phillip.melton@uwa.edu.au

Recommended Citation

Beaty, Kristine G.; Mosher, M. J.; Crawford, Michael H.; and Melton, Phillip, "Paternal Genetic Structure In Contemporary Mennonite Communities From The American Midwest" (2016). *Human Biology Open Access Pre-Prints*. 112.
http://digitalcommons.wayne.edu/humbiol_preprints/112

This Open Access Preprint is brought to you for free and open access by the WSU Press at DigitalCommons@WayneState. It has been accepted for inclusion in Human Biology Open Access Pre-Prints by an authorized administrator of DigitalCommons@WayneState.

Paternal genetic structure in contemporary Mennonite communities from the
American Midwest

Kristie Beaty¹, MJ Mosher², Michael H. Crawford³, Phillip Melton⁴

¹*University of Kansas*, ²*Western Washington University*, ³*University of Kansas Main Campus*, ⁴*University of Western Australia*.

Corresponding author: Phillip E. Melton, Ph.D Curtin/UWA Centre for Genetic
Origins of Health and Disease University of Western Australia 35 Stirling HWY
(M409) Crawley, WA, 6009 Australia Tel: 61 (8) 9224-0367 e-mail:
phillip.melton@uwa.edu.au

Keywords: Anabaptist, Y-Chromosome, Mennonites, Kansas, Nebraska, Population
Structure, Anthropological Genetics

Running header: Y-chromosome variation in Mennonites.

Issue: 88.2

Abstract:

Over the last 35 years, researchers from the Laboratory of Biological Anthropology at the University of Kansas have been working with Mennonite communities to better understand evolutionary patterns of fission-fusion in relationship to their genetic history and population structure. In this study, short tandem repeat (STR) markers from the non-recombining region of the Y-chromosome (NRY) were used to provide increased resolution of the molecular population structure for these groups.

This NRY is known to be informative for determining paternal genetic ancestral patterns in recently derived human populations. Mennonites represent a branch of the Anabaptist movement that began in northern and central Europe in the 16th century and maintain a well-documented migration and genealogical history. Provided this historical information, we investigated the genetic relationship of 15 NRY STR loci within five Mennonite communities from Kansas (Goessel, Lone Tree, Garden View, and Meridian) and Nebraska (Henderson). We sought to determine if patterns of fission/fusion along familial lines persisted with paternal genetic information as evidenced through other classical genetic polymorphisms and molecular markers. NRY haplotype information was obtained for 94 individuals and genetic variation were analysed and compared across the five study populations and comparative Anabaptist and European populations. NRY haplogroups were assigned using a Bayesian allele frequency approach using 14 STR loci. A total of 92 NRY haplotypes were detected, with none shared across these communities. The most prevalent NRY haplogroup was R1b, which occurred in 56% of the entire sample. Eight additional NRY haplogroups (E1b1b, G2a, I1, I2, J2a1, L, Q, and R1a) were detected in smaller frequencies. In contrast to mtDNA, principal component analysis of NRY data displayed no patterns of population subdivision of these communities into congregations. These NRY genetic profiles provide additional information regarding the recent migratory history of Mennonite communities and provide additional evidence for the fission along paternal lines after migration to the United States.

Modern Mennonite populations have gone through numerous historical migrations, with some communities settling in the Midwestern region of the United States. These

migrations are well documented both with historic and genealogical information and provide a unique opportunity for applying anthropological genetic approaches to examine Anabaptist population structure. Over the last 250 years, these Mennonite groups have inhabited three distinct geographic regions (Western Europe, Ukraine, and the United States). Their experiences at these three locations helped establish a unique cultural identity and a strong sense of shared community, particularly in the Ukraine, where they lived isolated from neighbouring Russian and non-Mennonite German settlers (Urry 1989; Stevenson and Everson 2000). Subsequent fissions and migration of these Mennonite groups were also impacted by schisms resulting from differences in religious ideology. Mennonite congregations inhabiting the Midwestern United States can be divided into three independent congregations, based on shared religious tenets within the Mennonite religious framework: (1) Alexanderwohl; (2) Holdeman; and (3) Old Colony. All three of these congregations have distinct demographic histories, and a number of them have been previously investigated using both classical (Comuzzie and Crawford 1990; Crawford and Rogers 1982; Crawford et al. 1989; Martin et al. 1996; Rogers 1984) and molecular genetic markers (Demarchi et al. 2005; Melton et al. 2010).

The Anabaptist movement started soon after the Reformation and is characterized by shared religious beliefs in adult baptism, separation of church and state, and pacifism. These groups represented the far left of the Reformation movement and arose in Switzerland, Germany, and the Netherlands (Rogers and Rogers, 2000). Anabaptist groups in these three regions were each associated with a charismatic leader and include: 1) Mennonite, followers of Menno Simons originating in northern Europe and the Netherlands; 2) Amish, followers of Jacob Amman, formed

in Switzerland and southern Germany; and 3) Hutterites, followers of Jacob Hutter, formed in Austria. After these groups formed, a number of small-scale rebellions broke out. Subsequently, local authorities began to view Anabaptists as a threat to social order, resulting in their persecution. These Anabaptist groups were forced to migrate either to underdeveloped areas of Eastern Europe or the Americas.

Anabaptist groups that migrated to the United States belong primarily to three distinct groups: 1) Swiss-south German groups, including the Amish; 2) Prussian Mennonites; and 3) Austrian Hutterites. Each of these Anabaptist groups have different cultural histories that may be reflected in their population structure. These ethnohistoric events are revealing and can be assessed in terms of their biological impact.

A brief overview of the migration history of these Mennonite communities is as follows. Dutch and German Mennonite refugees immigrated to Polish-controlled areas around Danzig (modern day Gdańsk, Poland), and in 1699, eighteen families formed the Przechowa church (L. Rogers and Rogers 2000; Krahn and Penner 2011). The population increased in size and maintained meticulous genealogical records. In 1821, all but seven families of the congregation moved to Russia and settled in the Ukraine near the Molotschna River. This congregation adopted the name Alexanderwohl, in honor of the Russian czar. Subsequent changes in economic conditions, shifts in Russian governmental policies concerning military exemptions, and internal subdivisions of these groups caused the Alexanderwohl Mennonites to migrate to the United States in 1874 (L. Rogers and Rogers 2000; Krahn and Penner 2011; Melton 2012). Upon arrival in the United States, the Alexanderwohl group split into two separate divisions (Goessel and Henderson). One group settled west of

Lincoln, Nebraska, near present-day Henderson. The other group settled in Kansas, 40 miles north of Wichita. A separate Kansas Mennonite congregation, founded in Ohio in 1858 by John Holdeman is representative of the Church of God in Christ Mennonites (Meridian, Garden View, Lone Tree). This congregation is considered a heterogeneous group composed of Pennsylvanian Dutch and Germans mixed with a large Mennonite immigrant populations from southern Russia (Crawford et al. 1989). The Meridian Holdeman Mennonite community further split after the 1980s into the congregations of Garden View, and Lone Tree.

A number of studies have investigated the genetic history of Mennonites using classical genetic polymorphisms (Crawford and Rogers, 1982; Rogers, 1984; Crawford et al 1989; Comuzzie and Crawford, 1990; Martin et al. 1996). This research has included blood group systems, serum proteins, and immunoglobulins (Crawford et al. 1989; Martin et al. 1996). More recent genetic studies on these Mennonite communities have focussed on molecular markers using Apolipoproteins (Demarchi et al. 2005) and mitochondrial DNA (mtDNA) diversity (Melton et al. 2010). This previous genetic research has demonstrated a fission-fusion pattern characterizing the recent evolutionary history and that these new Mennonite communities fission along familial lines (Crawford et al. 1989; Martin et al. 1996; Crawford 2005; Demarchi et al. 2005, Melton et al. 2010). Recent research on mtDNA in the Mennonite communities suggests that molecular genetic data provided a more accurate depiction of Anabaptist history than previously determined through classical genetic markers (Melton et al. 2010). However, mtDNA does not provide a complete genomic profile of a population and additional evidence from other

markers, including those within the non-recombining region of the Y-chromosome (NRY), is warranted.

Recently, anthropological genetic studies applied uniparental molecular genetic markers for examining the biological consequences resulting from migration of the different Anabaptist populations (Pollin et al. 2007; Melton et al. 2010; Pichler 2010). These Anabaptist groups (Amish, Hutterites, and Mennonites) have experienced dynamic histories characterized by several demographic events, which have contributed to their unique genetic structure (Martin et al. 1996; Crawford 2000; Melton et al. 2010; Melton 2012). However, to date few studies have focussed on the paternal contribution in these Anabaptist communities by examining NRY polymorphisms (Pollin et al. 2007; Pichler et al. 2010). Short tandem repeat (STR) markers from the NRY are known to be informative for determining paternal genetic ancestral patterns in recently derived human populations and examination of these polymorphisms provides additional insight into their genetic history. Two previous studies have investigated the NRY in Anabaptist communities. Pollin et al. (2007) studied NRY variation in the Amish population, and found a high correlation between their male genetic lineages and genealogical information based on surname analysis. Picher et al. (2010) investigated NRY variation in the Hutterite population and compared them to an Austrian population from South Tyrol, and found that this population demonstrated a unique genetic profile related to central and eastern European population. However, these studies did not compare Anabaptist populations to each other.

In the present study, we characterize NRY diversity within and between these distinct Midwestern Mennonite communities and assess their biological relationship with other Anabaptist and European populations. Our research aims for this article are to 1) determine the paternal genetic relationship between five Mennonite communities using NRY polymorphisms; 2) investigate the paternal biological relationship among two different Mennonite congregations and other European populations; and 3) determine if paternal population subdivision within these communities demonstrates patterns of fission-fusion as previously reported for classical genetic polymorphisms, immunoglobulins, and molecular markers.

MATERIALS AND METHODS

Population samples

We examined five Mennonite communities inhabiting Kansas and Nebraska subdivided into two major congregations: (1) Alexanderwohl, which includes the communities of Goessel, Kansas, and Henderson, Nebraska; and (2) Holdeman, which includes the Kansas communities of Meridian, Lone Tree, and Garden View. This study included 94 male participants, with samples collected as part of longitudinal multidisciplinary study of Midwestern Mennonite communities in the United States. Kansas samples were collected by researchers from the Laboratory of Biological Anthropology, University of Kansas in 2004, as described previously (Demarchi et al. 2005). Nebraska samples were collected in 1981 as part of a study of biological aging (Crawford 2000a). Human Ethics approval was approved by the

University of Kansas and signed informed consent was obtained for all participants in both studies.

Collection of blood samples and DNA extraction were performed as previously described (Melton et al. 2010). The 94 male samples used in this analysis included 13 individuals from Goessel, 21 from Henderson, 25 from Meridian, 15 from Garden View, and 20 from Lone Tree. To avoid close relatives we only investigated male participants with different surnames, which were checked against pedigree information to ensure accuracy. Comparative NRY STR data for Hutterites (Pilcher et al. 2010), Old Order Amish (Pollin et al. 2007), and eight European populations (Poland, Sweden, Netherlands, Finland, Italy, Russia, Germany, and Switzerland) were collected from the literature (Table 1).

NRY Analysis

Male participants were characterized for 15 NRY (DYS456, DYS389I & II, DYS390, DYS458, DYS19, DYS385 a/b*, DYS393, DYS391, DYS439, DYS635, DYS392, YGATAH4, DYS437, DYS438, and DYS448) STRs. These fifteen STRs were analysed using the AmpFISTR YFiler kit from Applied Biosystems (Foster City, CA) and multiplexed for fragment analysis on an ABI3130 sequencer at the University of Kansas Natural History Museum DNA Sequencing Laboratory. NRY STRs were assigned using Peak Scanner Software.1.0 (Applied Biosystems, Foster City, CA). NRY haplogroups were assigned using a Bayesian allele frequency approach using the 15 most informative NRY STR loci (www.hprg.com/hapest5/).

Analytical Techniques

Intrapopulation Analysis. NRY STR allelic frequencies, number of haplotypes, and additional diversity indices based on Nei (1987) were analysed using Arlequin 3.5 (Excoffier and Lischer 2010). Haplogroup frequencies were computed based on inferred assigned haplogroups as described above.

Population Structure. Population structure in the five Mennonite congregations was tested using analysis of molecular variance (AMOVA) to identify partitions of variance based on NRY STR data, and was performed in Arlequin 3.5 (Excoffier et al. 1992; Excoffier and Lischer 2010). AMOVA was also performed on previously analysed mtDNA hypervariable region 1 sequence data (Melton et al. 2010). Initial analyses were performed separating the communities by congregation (Alexanderwohl and Holdeman). An additional AMOVA analysis was performed, with the Alexanderwohl communities of Goessel and Henderson placed in the first group, Meridian and Garden View made up the second group, and Lone Tree was treated as a third group following results from Melton et al. (2010). Additionally, Mantel tests (Mantel 1967) were performed to determine the correlation between genetic distances and geographic distances. Pairwise distances were computed for NRY STR data (Current study; Pollin et al. 2007; Pichler et al. 2010) and for mtDNA data (van der Walt et al. 2005; Melton et al. 2010; Pichler et al. 2010) in Arlequin 3.5. Geographic distance matrices were calculated in R 3.2 (<https://www.r-project.org/>) and Mantel tests examining the relationship of Anabaptists NRY STR distances with mtDNA distances, NRY STR distances with geography, and mtDNA

sequence distances with geography were performed using the *ade4* package in R 3.2 (Dray et al. 2007).

Interpopulational Analysis. Principal coordinate analysis (PCoA) was used to visualize the biological relationships among Mennonite congregations and comparative European populations. Given differences between published NRY STR datasets, a reduced set of 6 common loci (DYS19, DYS389I & II, DYS390, DYS391, DYS392, and DYS393) were used to construct NRY STR pairwise distances in Arlequin 3.5. These distances were used to construct two dimensional PCoA plots using the APE package of R 3.2 (Paradis et al. 2004). Plots were constructed to examine the relationship of Anabaptist groups and to determine the relationship of these groups to parts of Europe that were briefly home to these refugees based on the historical records.

RESULTS

NRY STR Variation: A total of 94 individuals were characterized in these five Mennonite congregations and 92 different haplotypes were identified (Table 2), with no haplotypes shared between communities. Garden View and Lone Tree were the only two congregations where a single haplotype was identified in two individuals with gene diversities (H) of 0.9905 and 0.9947. When the data set excluded loci not found in the comparative literature, the two Alexanderwohl congregations of Goessel and Henderson shared two R1b haplotypes. The Holdeman communities also shared R1b haplotypes among the various congregations, and Henderson shared two R1b haplotypes with Meridian.

Haplogroup Distribution: Several European haplogroups were identified including haplogroups R1b (56.3%), R1a and I2 (9.6%), E1b (6.4%), I1 and Q (5.3%), G2a (3.2%), J2 (2.1%), and haplogroup L and an unidentified haplotype (1.1%). The distribution of these haplogroups varied among the congregations (Figure 2). Haplogroups R1b (50-63.2%) and R1a (4-26.3%) were the only haplogroups found in all 5 congregations. The Alexanderwohl congregations of Goessel and Henderson both exhibited haplotypes belonging to haplogroups G2a and J2a. Goessel exhibited the most haplogroup diversity with 6 haplogroups (R1b (53.8%), I2a (15.4%), J2a, G2a, R1a and Q (7.7%)) represented in 13 individuals and the highest mean number of pairwise differences between haplotypes (11.4872). Meridian exhibited the second highest haplogroup diversity with at least 7 NRY haplogroups present (R1b (56%), E1b1 (12%), I1a and Q (4%), I2, L, R1a, and one unidentified haplotype (4%)) in 25 individuals and an average number of pairwise differences of 11.22. It was the only congregation to have haplogroup L present. Lone Tree exhibited the lowest haplogroup diversity, with 63.2% of the individuals belonging to haplogroup R1b, 26.3% belonging to haplogroup R1a, and 5.3% representing haplogroups E1b1a and I2a; whereas Garden View had four haplogroups represented (R1b (50%), I2 (35.7%) R1b and Q (7.1%)) and exhibited the lowest average number of pairwise differences between haplotypes, 9.8476.

Population structure: AMOVA results for NRY STR and mtDNA variation are shown in Table 3. As with mtDNA, the amount of variation seen among groups is

lower than seen within all populations, while the amount of variation within populations is high. The amount of variation seen within populations is higher in mtDNA *vs.* NRY STR data. The amount of NRY STR variation explained among communities within each grouping is lower (6.12% *vs.* 7.57%) when Lone Tree is treated as a separate group from the other Holdeman communities. When Mantel tests are applied to these data and geographic proximity, geography and NRY variation is negatively correlated with mtDNA variation (Table 4). This is particularly true of NRY STR and mtDNA distances ($r = -0.5531$), however neither of these results gave significant p values. There is a slight correlation ($r = 0.1283$) of geography by NRY STR distances, but these results are also non-significant.

Intrapopulation Analyses: PCoA was used to determine the relationship of the Mennonite congregations with other Anabaptists groups using a reduced number of loci. The PCoA of Mennonite communities plots 77.5% of the NRY variation on the first two axes and shows that these Mennonite congregations are more similar to each other than they are to either the Old Order Amish or Hutterite Anabaptist populations (Figure 3). However, this PCoA also shows that Mennonite communities do not cluster by their original congregations of Alexanderwohl and Holdeman, and the community of Garden View pulled the farthest from the Mennonite communities in the plot. There is no sharing of paternal haplotypes between the Mennonite communities based on an expanded set of 15 NRY STRs, but with a reduced loci set of six STRs, sharing of haplotypes does occur (Table 5.). All of the shared paternal haplotypes between the communities belong to the most common and widespread western European NRY haplogroup, R1b. The sharing of paternal haplotypes also includes 4 R1b haplotypes shared with the Old Order Amish, with at least one of

these haplotypes found in each of the Mennonite communities. However, no Mennonite communities shared haplotypes with Hutterite populations included in this analysis, and this group is the furthest outlier within this plot.

A second PCoA plot was constructed comparing Anabaptists groups to other European populations using the reduced loci set of 6 NRY STRs and represents 46.2% of the observed NRY STR variation (Figure 4). The Mennonite communities plot nearest to Swiss, Italian, Dutch, German and Western Russian populations. All Mennonite communities share reduced haplotypes with Swiss and German groups, six of the 94 Mennonite paternal haplotypes are shared with the Dutch, and five haplotypes are shared with South Tyrol group from the Italian/Austrian border. The Hutterites are located to the far right of the plot, nearest to Western Russia and Swedish populations.

DISCUSSION

Population genetic studies examining religious isolates have long been known to be informative for the study of rare genetic disorders, due to their unique population structure (Pollin et al 2007; Pichler et al. 2010). However, very few studies have examined molecular genetic data to understand the diaspora of Anabaptist population following the Reformation. The primary focus for this current study was to determine the paternal genetic relationship between five Mennonite communities using NRY STR data, and to determine if these data support the history of fission-fusion that have occurred in these groups. Despite a shared history originating after the Reformation, there is no sharing of paternal haplotypes between Mennonite communities. The high level of paternal haplotype diversity seen in this sample may

be explained by the small male sample size, the exclusion of individuals with the same surname within each community, as well as the design of the Yfiler kit that is utilized in forensics to distinguish at an individual level. AMOVA analysis revealed that most of the paternal genetic variation observed in these groups is found within populations. This is not unexpected as NRY diversity among populations tended to be higher than that seen in mtDNA studies (Jorde et al. 2000; Pereira et al. 2001). NRY haplotypes tend to be more geographically specific due to differential genetic contributions of males versus females and kin migration (Mielke and Fix 2006). Despite this, some sharing of haplotypes may be expected due to the common origin of Mennonite groups. The PCoA plot of Anabaptists (Figure 3) populations demonstrates that Mennonite communities, while more closely related to each other than they are to other Anabaptist groups (Hutterites, Old Order Amish), do not cluster by congregation. These results may first appear to imply that belonging to a specific Mennonite congregation is not indicative of the NRY variation represented in this study. However, the history of these communities can be used to explain the distribution of NRY haplotypes and haplogroups seen in these Mennonite congregations.

As previously stated, the Alexanderwohl congregation migrated from the Ukraine to the United States in 1874. Upon arrival, this Alexanderwohl congregation split into two communities Henderson, Nebraska and Goessel, Kansas. This split was caused by differences in religious ideology, the availability of resources and additional economic factors (Rogers and Rogers 2000). While there is an absence of paternal haplotype sharing between these two settlements, there is evidence of a shared genetic ancestry. First, both communities demonstrate high frequencies of

haplogroup R1b (Goessel 53.8%, Henderson 52.4%), low frequencies of haplogroup R1a (7.7% and 4.8% respectively), and are the only Mennonite communities with Y haplogroups J2 and G2a. Furthermore, while there was no sharing of 15 loci NRY STR haplotypes, there were two 6 loci R1b haplotypes shared between the communities. There are also distinct differences in the paternal haplogroup distribution between these communities. The community of Goessel had 2 NRY haplogroups, I2a (15.4%) and Q (7.7%), that are not found in the Henderson community, while the Henderson sample had individuals belonging to haplogroup E1b1 (9.5%) These differences support the idea that related individuals tended to remain together (i.e. kin migration) when fission occurred between communities, and this is more pronounced occurring along related male lineages.

The Holdeman communities of Kansas are the descendants of a heterogeneous group of Pennsylvania Dutch and Germans that came to Ohio in 1858 and mixed with Kleine Gemeinde Mennonites that migrated from southern Russia in 1874 (Hiebert 1989). This heterogeneity is evident in the NRY STR data, as the original settlement of Meridian has the highest haplogroup diversity when compared to the other examined Mennonite communities. However, when these three Holdeman communities are compared to one another they share no NRY STR haplotypes and have very distinct paternal haplogroup compositions. Both Garden View and Lone Tree split from Meridian recently, and the result of this founder effect can be seen in the different distributions of non-R1b haplogroups. Garden View and Lone Tree contain NRY haplogroups I2 and R1a, but in Garden View the frequency of these haplogroups is 35.7% and 7.1%, whereas in Lone Tree the frequencies are at 5.3% and 26.3%. Both communities have haplogroups that were not represented in the

other, with the presence of haplogroup Q (7.1%) in Garden View and haplogroup E1b1 (5.3%) in Lone Tree. As with the Alexanderwohl congregation, this distribution of haplogroups can be the result of more related male individuals staying together when the group formed new communities away from Meridian.

The results from the NRY STR data from both Mennonite congregations indicate the movement of male lineages to new settlements in related groups of men, or kin-structured migration (Mielke and Fix 2006). Kin-structured migration, patrilocal residence patterns, and the STR kit design can explain why no Mennonite NRY STR haplotypes are shared among these five Mennonite communities. Melton et al. (2012) also noted that in these Mennonite communities, only 17 of the 87 surnames were found in more than one community, a result similar to an isonomy study of Mennonites by Rogers (1985). As surnames and NRY markers are both passed through paternal lineages in Western European societies, this unequal distribution of surnames and NRY haplotypes provides further evidence for kin structured fission in Mennonite populations.

Examination of the paternal biological relationship among the Mennonite congregations and other European populations provides for additional analysis regarding the genetic history of these communities. The unrest following the Reformation led to the dispersal and splintering of Anabaptists groups. Mennonites from the Netherlands and Germany migrated to Prussia between 1527 and 1539. Later unrest led Mennonite communities to flee to Eastern Europe, into Poland, in 1699, and later to the Ukraine (Rogers and Rogers 2000). Despite having inhabited areas of Eastern Europe, Mennonites have a typical Western/Northern European

genetic makeup, with high frequencies of R1b (56.4%), and a lower frequency of haplogroups I2 and R1a (9.6%). All Mennonite communities shared 6 loci haplotypes with Old Order Amish, Sweden, Switzerland, and Germany. At least three communities shared haplotypes with the Dutch, Western Russia, Finland, and Italy, indicating a primary affinity with Northern and Central European countries. This relationship is further illustrated in the PCoA plot (Figure 4), where Mennonite groups are most closely associated with Switzerland, Germany, Italy and the Dutch. Despite this, the Mennonite groups are distinct from their European source populations. Similar to studies of other Anabaptists groups (Pichler et al. 2010; Pollin et al. 2007) have shown that as these groups migrated from one region to another, there was relative isolation followed by periods of admixture. This movement and splintering of the Mennonite communities as a response to unrest led to bottlenecks in the various groups, followed by periods of genetic isolation from their host communities, and later admixture with other Mennonite groups.

A final aim of this study was to determine if the paternal population subdivision within these communities demonstrates patterns of fission-fusion seen in previous studies and to determine if molecular markers provided better subdivision between communities and congregations. Early studies of blood group polymorphism, serum proteins and immunoglobulins showed mixed results. A study by Crawford et al. (1989) using 19 polymorphic classical markers showed a common history of the Alexanderwohl communities, with more similarities between these two communities than between either with the Holdeman community of Meridian. When the study was expanded to include 44 allele frequencies and 15 classical genetic markers a different result emerged, with the Kansas communities of Meridian and Goessel appearing

more similar to each other than they were to the Nebraska community of Henderson. This result countered that from the historical record but could be explained through kin-structured migration (Crawford, 2000). The Nebraska community was found to have a unique RHR haplotype that distinguished it from the Kansas communities, indicating that founder effect may have contributed to the observed differences. These results are more similar to that seen in the current study, where clustering by congregation did not occur. Martin et al. (1996) utilized variation in immunoglobulins, genetic markers that allow for higher resolution and more population specific than blood group polymorphisms, to better understand the genetic relationship of three Mennonite communities (Goessel, Henderson, and Meridian). These studies identified small differences in haplotype frequencies among Mennonite communities, variation that could be explained by non-random fission along familial lines.

The mixed results of early molecular markers illustrated the need to apply higher resolution molecular genetic techniques to better understand the patterns of fission-fusion seen in the Mennonites. In 2010, Melton et al. used mtDNA HVS-I sequence variation in these same five Mennonite communities to provide higher resolution of the maternal genetic variation to and provide a female history to Mennonite migration. As with the earlier study of classical polymorphisms (Crawford et al. 1989), mtDNA variation could be attributed to genetic ancestry based on congregational affiliation. The study suggested that maternal variation split along familial lines, but that these lineages were more related to one another within congregations than some of the other variants examined. Again, these findings are different from those seen in the present study of NRY markers for these groups, yet

some similarities remain. Both sets of molecular markers show a high genetic diversity values indicative of low inbreeding, despite the tendency to select mates that were also Mennonites. This is supported by marriage records collected by Stevenson and Everson (2000) that found that Mennonite marriages outside of congregational affiliations were relatively high during the 1700s – early 1800s, with mates found from other congregations 20 – 50 percent of the time. Once in Russia, these rates dropped below 2 percent. However, reproductive isolation of Mennonite congregations has reduced since the 1930s, with mate selection occurring outside of the congregation and sometimes outside of Mennonite communities (Stevenson and Everson 2000). The differences of clustering when comparing Mennonite mtDNA sequences and NRY STRs may be explained if fission occurred along male and female lines when communities were formed in the United States, but that women might have been more likely to move between other communities within the same congregations, a patrilocal residence pattern common among Anabaptist populations. These results demonstrate the utility in using uniparental molecular markers, as they give a more accurate picture of the history of the Mennonites of the Midwest that is consistent with documented historical and genealogical records.

CONCLUSION

This study investigated NRY STR markers in five Mennonite communities and related populations to examine kin-structured migration in paternal lineages for these groups. In contrast to evidence from mtDNA and some classical genetic polymorphisms, NRY variation does not segregate these Mennonite communities by their associated congregation but suggests that male relatives migrate and settle together when new communities are formed. These results demonstrate clear

evidence that Mennonite groups share genetic affinities with Central and Northern Europe, areas in which they originated. However, these groups underwent a series of fission and fusion events in their recent evolutionary history, resulting in the current observed NRY variation, where fissions appear to have occurred recently along paternal lines after migration to the United States. Provided this short time frame and the use of uniparental markers that are susceptible to kin-structured migration, additional evidence from genome-wide autosomal markers may provide a more balanced examination of Mennonite genetic ancestry.

Acknowledgements This research was sponsored by an AAPA Professional Development Award, Kansas Attorney General Settlement grant, and the National Institute of Aging. Research by PEM is supported in part by funding from The Royal Perth Hospital Medical Research Foundation. We thank all Mennonites individuals who participated in this study.

Received 9 December 2015; revision uploaded 17 July 2016.

REFERENCES

- Allen, G. 1988. Random genetic drift inferred from surnames in Old Colony Mennonites. *Hum. Genet.* 60:639-653.
- Allen, G., and C. W. Redekop. 1967. Individual differences in survival and reproduction among Old Colony Mennonites in Mexico: Progress to October 1966. *Eugen. Quart.* 14:103-111.
- Allen, G., and C. W. Redekop. 1987. Old Colony Mennonites in Mexico: migration and inbreeding. *Soc. Biol.* 34:166-179.
- Comuzzie, A. G., and M. H. Crawford. 1990. Biochemical heterozygosity and morphological variability: interpopulational versus intrapopulational analyses. *Hum. Biol.* 62(1):101-112.

Crawford, M. H. 2005. Genetics of biological aging in Mennonites of Midwestern United States. *Przegląd Antropologiczny* 68:3-18.

Crawford, M. H. 2000. Genetic structure of Mennonite populations. In *Different Seasons: Biological Aging among the Mennonites of the Midwestern United States*. M.H. Crawford, ed. University of Kansas, 31-40.

Crawford, M. H., D. D. Dykes, and H. F. Polesky. 1989. Genetic structure of Mennonite populations of Kansas and Nebraska. *Hum. Biol.* 61:493-514.

Crawford, M. H., and L. Rogers. 1982. Population genetics models in the study of aging and longevity in a Mennonite community. *Soc. Sci. Med.* 16:149-153.

Demarchi, D., M. J. Mosher, and M. H. Crawford. 2005. Apolipoproteins (apoproteins) and LPL variation in Mennonite populations of Kansas and Nebraska. *Am. J. Hum. Biol.* 17: 593-600.

Dray, S., and A. B. Dufour. 2007. The ade4 package: implementing the duality diagram for ecologists. *J. Stat. Softw.* 22(4): 1-20. Version 1.7-2, April 14, 2015. <http://pbil.univ-lyon1.fr/ADE-4>

Dupanloup, I., S. Schneider, and L. Excoffier. 2002. A simulated annealing approach to define the genetic structure of populations. *Mol. Ecol.* 11(12):2571–2581.

Excoffier, L., P. E. Smouse, and J. M. Quattro. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131: 479-491.

Excoffier, L., and H. E. L. Lischer. 2010. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10: 564-567. <http://cmpg.unibe.ch/software/arlequin35/>

Hiebert, P. G., and C. Hiebert. 1989. Church of God in Christ, Mennonite (CGC). *Global Anabaptist Mennonite Encyclopaedia Online*. Retrieved 29 November 2015, from [http://gameo.org/index.php?title=Church_of_God_in_Christ,_Mennonite_\(CGC\)&oldid=120959](http://gameo.org/index.php?title=Church_of_God_in_Christ,_Mennonite_(CGC)&oldid=120959).

Jorde, L. B., W. S. Watkins, M. J. Bamshad et al. 2000. The distribution of human genetic diversity: a comparison of mitochondrial, autosomal and Y-chromosome data. *Amer. J. Hum. Genet.* 66(3): 979-988.

Karlsson, A. O., T. Wallerström., A. Götherström et al. 2006. Y-chromosome diversity in Sweden—a long-time perspective. *Eur. J. Hum. Genet.* 14(8):963-970.

Kayser, M., M. Krawczak, L. Excoffier et al. 2001. An extensive analysis of Y chromosomal microsatellite haplotypes in globally dispersed human populations. *Am. J. Hum. Genet.* 68:990-1018.

Krahn, C., and G. Penner. 2011. Alexanderwohl (Molotschna Mennonite Settlement, Zaporizhia Oblast, Ukraine). *Global Anabaptist Mennonite Encyclopedia Online*. Retrieved 1 December 2015, from [http://gameo.org/index.php?title=Alexanderwohl_\(Molotschna_Mennonite_Settlement,_Zaporizhia_Oblast,_Ukraine\)&oldid=132408](http://gameo.org/index.php?title=Alexanderwohl_(Molotschna_Mennonite_Settlement,_Zaporizhia_Oblast,_Ukraine)&oldid=132408).

Lappalainen, T., S. Koivumäki, E. Salmela et al. 2006. Regional differences among the Finns: a Y-chromosomal perspective. *Gene* 376(2):207-215.

Mantel, N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Res.* 27:209-220.

Martin, K, J. C. Stevenson, M. H. Crawford et al. 1996. Immunoglobulin haplotype frequencies in Anabaptist population samples: Kansas and Nebraska Mennonites and Indiana Amish. *Hum. Biol.* 68:45-62.

Melton, P. E. 2012. Mennonite migrations: genetic and demographic consequences. In *Causes and Consequences of Human Migration: an Evolutionary Perspective*. Crawford, M.H. and B. C. Campbell eds. Cambridge University Press, 299-316.

Melton, P. E., M. J. Mosher, R. Rubicz et al. 2010. MtDNA Diversity in Midwestern Mennonites. *Hum. Biol.* 82(3):267-289.

Mielke, J. H., and A. G. Fix. 2006. The confluence of anthropological genetics and anthropological demography. In *Anthropological Genetics: Theory, Methods and Applications*. Crawford, M.H. and B. C. Campbell eds. Cambridge University Press, 112-140.

Myers, N. M., S. Roostsi, A. A. Lin et al. 2011. A major Y-chromosome haplogroup R1b Holocene era founder effect in Central and Western Europe. *Eur. J. Hum. Genet.* 19:95-101.

Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20: 289-290. Version 3.3, May 28, 2015. <http://ape-package.ird.fr/>

Pereira, L., I. Dupanloup, Z. H. Rosser et al. 2001. Y-chromosome mismatch distributions in Europe. *Mol. Biol. Evol.* 18(7):1259-1271.

Pichler, I., C. Fuchsberger, C. Platzer et al. 2010. Drawing the history of the Hutterite population on the genetic landscape: inference from Y-chromosome and mtDNA genotypes. *Eur. J. Hum. Genet.* 18(4):509.

Pichler, I., J. C. Mueller, S. A. Stefanov et al. 2009. Genetic structure in contemporary south Tyrolean isolated populations revealed by analysis of Y-chromosome, mtDNA, and Alu polymorphisms. *Hum. Biol.* 81:875-898.

- Pollin, T. I., D. J. McBride, R. Agarwala, et al. 2007. Investigations of the Y chromosome, male founder structure and YSTR mutation rates in the Old Order Amish. *Hum. Hered.* 65(2):91-104.
- Roewer, L., S. Willuweit, C. Krüger et al. 2008. Analysis of Y chromosome STR haplotypes in the European part of Russia reveals high diversities but non-significant genetic distances between populations. *Int. J. Legal Med.*, 122(3):219-223.
- Rogers, L. A. 1984. Phylogenetic identification of a religious isolate and the measurement of inbreeding. Ph.D. dissertation, University of Kansas, Lawrence.
- Rogers, L., and R. A. Rogers. 2000. Mennonite history with special reference to Alexanderwohl and related congregations in Kansas and Nebraska. In *Different Seasons: Biological Aging among the Mennonites of the Midwestern United States*, University of Kansas, 7-18.
- Slatkin, M. 1995. A measure of population subdivision based on microsatellite allele frequencies. *Genetics* 139:457-462.
- Stevenson, J., and P. Everson. 2000. Historical demography of Mennonite populations. In *Different Seasons: Biological Aging among the Mennonites of the Midwestern United States*, University of Kansas 21, 19-30.
- Trynova, E. G., T. N. Tsitovich, E. Y. Vylegzhanina et al. 2011. Presentation of 17 Y-chromosomal STRs in the population of the Sverdlovsk region. *Forensic Sci. Int.-Gen.* 5(3):e101-e104.
- Urry, J. 1989. Mennonite economic development in the Russian mirror. In *Mennonites in Russia, 1788-1988: Essays in Honour of Gerhard Lohrenz*, 99-126.
- van der Walt, J. M., W. K. Scott, S. Slifer et al. 2005. Maternal lineages and Alzheimer disease risk in the Old Order Amish. *Hum Genet.* 118(1):115–122

FIGURE LEGENDS




Figure 1. Mennonite Communities of the Midwest reprinted with permission of Melton et al. (2010).




Figure 2. Map of Mennonite congregations that participated in this study with the frequencies of predicted haplogroups found in each.

Anabaptists PCoA Ordination




Figure 3. Principal coordinate analysis of Anabaptist groups including the five Mennonite communities, Hutterites, and Old Order Amish.




Figure 4. Principal coordinate analysis of Anabaptist groups with other European populations.

Group/Country	Subgroup	n	Source (Y STRs/mtDNA)	latitude	longitude
Mennonite	Goessel	13	Current study/Melton et al. 2010	38.2464	-97.3489
	Gardenview	15		38.0819	-97.5137
	Henderson	21		40.7797	-97.8123
	Lonetree	20		38.3054	-97.5464
	Meridian	25		38.2275	-97.4165
Hutterites	South Tyrol	75	Pichler et al. 2009/Pichler et al. 2010	46.7341	11.2888
Old Order Amish	Lancaster County, PA	732	Pollin et al. 2007/van der Walt et al. 2005	40.0467	-76.1784
Sweden	Blekinge/Kristianstad	41	Karlsson et al. 2006	na	na
	Gotland	40		na	na
	Swedish Saami	38		na	na
	Skaraborg	45		na	na
	Uppsala	55		na	na
	Värmland	42		na	na
	Västerbotten	41		na	na
	Österbotten	40		na	na
Östergötland/Jönköping	41	na	na		
Western Russia	Archangelskaja	42	Roewer et al. 2008	na	na
	Brianskaja	43		na	na
	Iwanovskaja	40		na	na
	Lipezkaja	47		na	na
	Nowgorodskaja	40		na	na
	Orlovskaja	42		na	na
	Penzenskaja	81		na	na
	Rjazanskaja	36		na	na
	Smolenskaja	42		na	na
	Tambovskaja	48		na	na
	Tverskaja	42		na	na
	Vologodskaja	40	na	na	
	Sverdlovsk	32	Trynova et al. 2011	na	na
Switzerland	Switzerland	64	Kayser et al. 2001	na	na
German	German	88		na	na
Dutch	Dutch	88		na	na
Poland	Poland	208	Rebala & Szczerkowska 2004	na	na
Italy	South Tyrol	227	Pichler et al. 2010	na	na
East Finland	Northern Karelia	22	Lappalainen et al. 2006	na	na
	Northern Ostrobothnia	129		na	na
	Northern Savo	107		na	na
	Southern Karelia	48		na	na
West Finland	Hame	49		na	na
	South-Western Finland	50		na	na
	Southern Ostrobothnia	58		na	na
	Swedish Spk Ostrobothnia	25		na	na
	Satakunta	47		na	na

Table 1. Populations used in this study

Table 2. NRY Haplogroups Identified in the Five Mennonite Communities

Group	Haplogroup (Probability)	Y STR Loci															# of Haplotypes shared (Reduced loci set in grey)																					
		DYS19	DYS3891	DYS3891I	DYS390	DYS391	DYS392	DYS393	DYS385	DYS437	DYS438	DYS439	DYS448	DYS456	DYS458	DYS635	Y-GATA_H4	Goessel	Henderson	Garden View	Lonetree	Meridian	Amish	Hutterites	W Russia	Sweden	Switzerland	E Finland	W Finland	Germany	Poland	Italy	Dutch					
Goessel	G2a(99.9%)	15	12	27	21	12	13	14	14,15	16	10	11	22	14	17	25	12																					
	I2a(58.6%)	14	14	26	26	11	10	13	12,15	14	10	10	21	17	18	25	12																					
	I2b1(70.9%)	17	14	28	24	10	12	14	14,15	15	12	10	20	15	17	22	11																					
	J2a1x (90.8%)	13	14	30	20	11	10	13	12,15	14	9	12	21	15	16	23	12																					
	Q(93.7%)	13	13	27	21	11	10	13	14,15	15	13	13	20	17	16	22	12																					
	R1a(93.7%)	16	13	30	25	11	11	14	12,15	16	11	11	20	15	17	22	13								3				1		1							
	R1b(100%)	14	15	33	26	12	13	13	12,15	16	13	12	19	17	18	24	11																					
	R1b(100%)	14	14	25	25	11	9	13	12,15	15	13	12	19	17	16	24	12																					
	R1b(100%)	14	14	31	24	11	13	13	12,15	15	13	12	19	15	16	23	12		1							3	1											
	R1b(100%)	14	14	32	25	11	13	13	12,15	15	13	12	19	17	16	26	12																					
	R1b(100%)	14	14	32	24	12	13	13	14,15	15	13	13	19	15	17	26	11		1																			
	R1b(83.8%)	15	14	26	26	12	11	13	11,12	14	12	13	19	16	17	24	12																					
R1b(97.2%)	14	11	33	26	10	10	12	14,15	14	13	12	20	18	16	24	13																						
Henderson	E1b1a(88.1%)	15	15	32	21	10	12	15	14,16	14	11	11	20	15	16	22	10																					
	E1b1b(93.1%)	16	15	33	24	10	12	13	15,17	14	11	11	20	16	17	22	10																					
	G2a(76.2%)	15	13	30	23	10	11	13	14,15	16	11	12	20	15	16	24	12																					
	G2a(83.7%)	15	14	31	20	10	11	14	12,15	17	13	10	20	17	17	23	13																					

	I1(68.8%)	14	13	30	23	10	11	13	12,15	16	11	11	20	15	16	24	11										
	I1(99%)	14	13	29	23	10	11	13	14,15	17	11	10	20	15	16	25	12	1		2	2		2	3			
	I1(99.4%)	14	13	30	23	12	11	13	14,15	16	11	11	20	16	15	24	10										
	J2a1h(98.1%)	15	14	30	24	10	11	12	15,17	14	10	11	21	16	14	24	12			2							
	Q(68.8%)	15	14	33	24	10	12	13	15,17	15	11	11	20	16	16	22	13										
	R1a(100%)	16	14	31	26	10	11	13	14,15	14	12	10	20	17	15	24	13										
	R1b(100%)	14	14	32	24	12	13	13	12,15	16	13	14	19	16	18	24	12	1									
	R1b(100%)	14	14	30	25	11	13	13	12,15	16	13	13	20	17	17	24	13		1								
	R1b(100%)	14	10	31	25	11	13	15	12,15	14	13	12	18	17	17	24	11										
	R1b(100%)	14	14	31	24	10	13	12	14,15	15	13	10	19	16	19	23	12										
	R1b(100%)	15	14	30	25	11	13	13	12,15	16	13	11	17	17	19	24	12										
	R1b(100%)	14	14	30	24	12	13	13	12,15	15	13	10	19	17	17	24	11								1		
	R1b(100%)	14	14	31	24	11	13	13	12,15	15	13	12	19	17	16	24	12	1		41		3	1				
	R1b(100%)	14	13	29	24	12	13	13	12,15	15	13	12	19	17	17	24	12			4							
	R1b(100%)	14	14	30	25	11	13	13	12,15	15	13	13	21	17	17	25	12		1							1	
	R1b(100%)	14	14	30	25	11	13	13	12,15	15	13	13	21	17	17	25	12		1							1	
	R1b(77.6%)	14	13	29	23	10	11	13	14,15	15	11	12	19	15	16	25	12		1			2	2		2	3	
	R1b(99.8%)	14	13	30	24	10	14	13	12,15	14	12	11	20	17	16	23	12									1	
Gardenview	I2a(74.7%)	17	14	29	25	11	8	13	12,15	15	13	9	19	16	16	24	8										
	I2a(77%)	15	14	26	24	11	13	13	13,16	15	10	12	20	15	16	24	12										
	I2a(86.6%)	15	13	31	24	12	16	13	12,15	15	11	15	21	15	19	24	12										
	I2a(93.5%)	14	13	30	24	10	10	13	12,15	15	10	13	18	16	17	24	12										
	I2a(95.9%)	14	14	28	24	11	11	13	12,15	14	8	12	20	16	17	24	12										

	Q(67.7%)	14	14	31	24	11	16	13	12,15	15	11	14	20	16	18	25	12											
	R1a(50.2%)	13	14	31	22	11	18	13	11,15	14	10	11	20	17	15	23	10											
	R1b(100%)	14	14	25	25	11	13	13	12,15	16	12	12	17	17	16	23	12											
	R1b(100%)	15	13	30	25	11	13	13	12,15	16	13	12	17	17	16	24	12											
	R1b(100%)	16	14	30	26	11	13	13	12,15	15	13	14	18	16	18	23	12											
	R1b(100%)	14	14	30	24	10	13	13	12,15	15	13	14	18	17	16	23	12		1	20	1	1	1	1	1			
	R1b(100%)	12	15	24	24	11	13	13	12,15	15	13	12	19	17	17	26	11											
	R1b(100%)	14	15	32	24	10	13	13	12,15	15	13	12	19	17	17	25	11											
	R1b(100%)	14	14	30	26	11	13	13	12,15	14	13	11	17	16	18	24	11	1		1								
	R1b(100%)	14	14	30	26	11	13	13	12,15	14	13	11	17	16	18	24	11	1		1								
Lonetree	E1b1b(94.2%)	13	13	30	25	10	12	13	12,16	15	8	13	21	18	17	24	11											
	I2a (88%)	15	13	29	24	10	11	13	10,15	17	13	12	20	15	15	21	11			3								
	R1a (100%)	16	14	32	26	10	11	13	11,19	15	12	11	19	17	15	24	12					1						
	R1a (100%)	16	15	32	26	10	8	13	12,16	14	12	11	20	17	15	24	13											
	R1a(100%)	16	14	32	26	10	11	13	12,15	14	12	11	19	17	15	23	12					1						
	R1a(100%)	16	13	31	26	10	11	13	13,15	14	12	11	20	16	15	24	13											
	R1a(93.1%)	16	14	32	25	10	18	13	21,25	14	11	13	20	16	15	24	11											
	R1b (99.4%)	14	14	24	24	11	15	13	12,15	16	13	13	23	16	16	24	13											
	R1b(100%)	14	13	30	24	10	13	13	11,12	14	13	13	19	16	16	23	11					2		4		1	1	1
	R1b(100%)	14	13	30	24	10	13	13	12,15	15	13	12	18	16	17	24	12					2		4		1	1	1
	R1b(100%)	14	14	30	24	10	13	13	12,15	14	12	10	19	17	18	23	11		1		20	1	1	1	1	1	1	
R1b(100%)	14	14	32	24	11	13	13	12,15	15	13	12	20	16	17	24	12												

	R1b(100%)	14	15	32	24	11	13	13	12,15	15	13	12	19	17	17	25	11	1										
	R1b(100%)	14	13	30	24	10	13	13	12,15	15	13	12	18	16	17	24	12	2		4		1		1	1			
	R1b(100%)	14	15	32	24	11	13	13	12,15	15	13	12	19	16	17	24	11	1										
	R1b(82.7%)	15	13	30	24	10	16	13	11,13	16	12	8	20	15	17	22	11											
	R1b(96.4%)	15	14	26	24	10	15	13	14,15	14	13	11	20	16	16	24	12											
	R1b(98.7%)	14	13	29	23	10	10	13	12,15	16	11	12	20	15	17	23	11											
	R1b(99.5%)	16	14	32	25	10	13	13	12,15	15	13	13	19	16	17	23	10											
	R1b(99.9%)	15	14	24	24	10	9	13	11,17	15	9	14	19	16	17	24	11											
Meridian	E1b1b (81.2%)	16	13	32	23	10	12	13	15,17	16	11	12	20	17	16	22	11											
	E1b1b (99.8%)	13	14	32	23	10	10	13	12,15	14	10	12	20	17	15	23	12											
	E1b1b(89.1%)	13	15	32	24	13	11	13	15,17	15	13	13	22	15	18	24	11											
	I1(72.6%)	14	13	29	23	10	12	12	12,15	16	12	11	20	14	15	24	11		1									
	I1(99.2%)	14	13	29	23	10	12	12	12,15	16	12	11	20	14	15	22	11		1									
	I2a(93.4%)	15	13	27	23	11	12	13	12,15	14	11	13	21	15	14	25	12											
	L(99.6%)	13	14	27	23	10	15	13	15,17	16	10	14	20	15	17	24	12											
	Q(76.7%)	13	13	31	23	12	14	13	12,15	14	10	13	22	15	18	24	13											
	Q(97%)	13	13	29	25	10	11	13	15,17	15	12	13	21	14	18	25	12											
	R1a(99.6%)	13	14	33	27	11	11	13	12,15	14	10	10	21	17	17	24	11											
	R1b (100%)	14	14	30	24	11	10	13	12,15	15	13	14	19	15	19	24	12											
	R1b(100%)	14	14	30	24	11	13	13	12,15	15	13	11	19	17	17	23	12		1		1	3	1		2	3	1	1
	R1b(100%)	15	14	31	26	11	13	13	12,15	16	13	12	18	17	17	23	11											
	R1b(100%)	14	14	30	25	11	13	13	12,15	15	12	12	20	15	17	23	13		2									
R1b(100%)	14	14	29	24	11	13	13	12,15	16	13	12	19	17	17	23	13			2						1			

R1b(100%)	14	13	26	20	10	12	13	11,13	14	13	14	20	16	16	23	12
R1b(100%)	13	11	32	20	13	12	14	12,15	15	12	12	20	15	16	25	12
R1b(100%)	14	14	30	26	11	10	13	12,15	14	13	11	18	15	18	23	11
R1b(100%)	14	14	30	26	11	13	13	12,15	16	13	11	20	15	18	23	11
R1b(100%)	14	14	30	24	11	13	13	12,15	15	13	15	20	15	19	24	12
R1b(84.0%)	15	12	29	20	12	13	13	12,15	15	13	13	20	16	16	22	12
R1b(96.2%)	12	14	31	23	11	14	13	12,15	15	11	14	20	17	16	25	12
R1b(99.8%)	13	14	33	21	12	13	13	12,15	15	13	14	20	15	16	25	12
R1b(99.8%)	16	13	31	21	11	13	12	12,15	14	13	14	20	16	16	24	11
unknown	16	13	30	21	12	13	13	12,15	13	12	14	20	14	16	23	12

2

1

1

3

1

2

3

1

1

Y STR

% of variation	Source of Variation	d.f.	Sum of squares	Variance components	% of variation
9.89	Among Groups	4	102.343	0.22122	11.29
7.57	Among pop. w/in Groups	2	7.416	0.12002	6.12
82.54	W/in Populations	816	1321.018	1.61889	82.59
	Total	822	1430.776	1.96013	

Fixation Indices	
FSC:	0.06902
FST:	0.17409
FCT:	0.11286

mtDNA

% of variation	Source of Variation	d.f.	Sum of squares	Variance components	% of variation
9.48	Among Groups	4	23.029	0.19586	10.12
1.25	Among pop. w/in Groups	2	3.028	-0.01224	-0.63
89.28	W/in Populations	111	194.351	1.75091	90.51
	Total	117	220.407	1.93453	

Fixation Indices	
FSC:	-0.00704
FST:	0.09492
FCT:	0.10125

*Lonetree separate group

Table 3. AMOVA Results for NRY STRs and mtDNA Sequences

Mantel test	r	p-value
Geograpy x mtDNA	-0.03549	0.5345
Y STR diversity x mtDNA	-0.55531	0.9932
Geography x Y STR diversity	0.128335	0.4427

Table 4. Mantel Tests of Geographic Proximity, mtDNA, and NRY STR Diversity

		Percent of Haplotypes Shared													
	Goessel	Gardenview	Henderson	Lonetree	Meridian	Old Order	Hutterites	Wrussia	Sweden	Switzerland	Efinland	Wfinland	Germany	Italy	Dutch
Goessel			9.52			5.60		0.55	0.52	0.67			0.31		
Gardenview				5.00	4.00	2.73			0.26	0.67	0.33		0.31		1.14
Henderson	15.38				4.00	6.15		0.55	1.04	1.34	0.98	1.31	0.31	0.44	1.14
Lonetree		6.67				2.73		0.55	1.31	0.67	0.33		0.62	0.44	2.27
Meridian		13.33	9.52			0.27		0.18	0.78	1.34		0.87	0.93	0.88	2.27
Old Order	7.69	6.67	9.52	5.00	4.00		21.33	2.57	12.27	18.12	0.98	3.49	10.56	3.96	7.95
Hutterites						21.86		8.44	18.28	13.42	5.88	9.61	9.01	4.41	3.41
Wrussia			14.29	0.05	4.00	23.22	45.33		51.96	26.85	25.82	41.92	41.61	11.01	17.05
Sweden	7.69	6.67	19.05	20.00	4.00	50.82	65.33	44.40		49.66	32.68	47.16	37.89	13.66	27.27
Switzerland	7.69	6.67	9.52	5.00	4.00	50.96	57.33	17.43	37.08		10.13	16.59	31.06	11.45	22.73
Efinland		6.67	14.29	5.00		25.96	62.67	31.56	43.60	19.46		41.48	15.22	4.41	6.82
Wfinland			9.52		4.00	25.27	62.67	29.91	49.09	20.81	32.03		24.53	6.61	11.36
Germany	7.69	6.67	4.76	20.00	4.00	46.99	80.00	54.86	43.08	51.01	12.09	24.45		18.06	46.59
Italy			4.76	15.00	4.00	69.81	86.67	18.90	40.47	37.58	8.17	17.47	29.19		21.59
Dutch		6.67	4.76	20.00	8.00	30.60	57.33	7.71	30.81	35.57	6.86	16.16	27.64	9.69	
Total hts	13	15	21	20	25	732	75	545	383	149	306	229	322	227	88

Table 5. Percentage of NRY Haplotypes Shared between Comparative Populations