

2009

Failure of *E. coli* bacteria to induce preterm delivery in the rat

Emmet Hirsch

NorthShore University HealthSystem, ehirsch@northshore.org

Yana Filipovich

NorthShore University HealthSystem, yfilipovich@netscape.net

Roberto Romero

Perinatology Research Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Department of Health and Human Services, prbchiefstaff@med.wayne.edu

Recommended Citation

Hirsch *et al.* *Journal of Negative Results in BioMedicine* 2009, **8**:1
doi:10.1186/1477-5751-8-1

Available at: <http://digitalcommons.wayne.edu/biomedcentral/135>

Research

Open Access

Failure of *E. coli* bacteria to induce preterm delivery in the rat

Emmet Hirsch*^{1,2}, Yana Filipovich¹ and Roberto Romero³

Address: ¹Department of Obstetrics and Gynecology, NorthShore University HealthSystem, Evanston, IL, USA, ²Department of Obstetrics and Gynecology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA and ³Perinatology Research Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Department of Health and Human Services, Bethesda, MD and Detroit, MI, USA

Email: Emmet Hirsch* - ehirsch@northshore.org; Yana Filipovich - yfilipovich@netscape.net; Roberto Romero - prbchiefstaff@med.wayne.edu

* Corresponding author

Published: 4 January 2009

Received: 8 August 2008

Journal of Negative Results in BioMedicine 2009, **8**:1 doi:10.1186/1477-5751-8-1

Accepted: 4 January 2009

This article is available from: <http://www.jnrnm.com/content/8/1/1>

© 2009 Hirsch et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: We sought to develop a model of bacterially induced preterm delivery in rats to parallel similar models in mice.

Methods: Female Sprague-Dawley rats on day 17 of gestation (normal term = 21–22 days) were inoculated into the uterus with either 2×10^9 – 7×10^{10} killed *E. coli* organisms, 1 – 4×10^8 live *E. coli* or sterile solution. These inoculations were made either via trans-cervical catheter or by direct intrauterine injection at laparotomy. Animals were then observed for delivery for variable periods up to term. Necropsies were performed and fetal viability was assessed.

Results: No rats delivered prematurely after bacterial exposure (27 animals observed for at least 48 hours), and all animals followed to term ($n = 3$) delivered live pups. No dams exhibited signs of systemic illness. There was a statistically significant but small negative effect of killed *E. coli* on fetal viability (100% of 80 fetuses from 6 control pregnancies and 93% of 182 fetuses from 14 bacterially-treated pregnancies were alive at necropsy, $p = 0.014$). Live bacteria had a larger effect on fetal viability, with only 64% of 14 fetuses, 47% of 28 fetuses and 32% of 31 fetuses surviving after trans-cervical administration of 7×10^7 , 2×10^8 and 4×10^8 *E. coli*, respectively.

Conclusion: Unlike mice, it has proven difficult to induce preterm labor in the rat using *E. coli* as a stimulating agent. The relevant literature is reviewed and hypotheses are offered to explain this phenomenon.

Background

Preterm birth is the major cause of neonatal morbidity and mortality in the developed world [1]. Infection within the gestational compartment is thought to be the primary cause of a large proportion of cases of preterm labor, accounting for as many as 50% or more of premature deliveries, especially at very early gestational ages. Given the complexity inherent in the processes of parturition and the obstacles to conducting properly controlled and

prospective studies in human subjects, animal models have proven helpful in developing an understanding of the physiology and pathophysiology of parturition [2]. Novel insights have been generated in animal models of infection-induced preterm labor in mice using a wide range of inflammatory stimuli from a large number of labs (small representative sample found in references [3–9]), including our own. Other well established models exist in rabbits [10–13] and non-human primates [14–16].

In order to capitalize on certain advantages afforded by the rat over the mouse (such as larger tissue samples and more easily conducted invasive monitoring), we sought to develop a model of infection-induced preterm labor in the rat. In this paper we report on the failure of this attempt, offer hypotheses for the causes of this failure and suggest possible directions for future study. It is notable that although there are abundant studies of rat parturition induced by progesterone antagonism [17-19] only two groups have reported preterm delivery with an infectious/inflammatory stimulus in rats in a total of three publications [20-22].

Methods

Rats

All procedures involving animals were approved by the Institutional Animal Care and Use Committee of North-Shore University HealthSystem. Timed pregnant Sprague-Dawley rats (254 – 380 gm) were obtained from Harlan (Indianapolis, IN) on day 15 of pregnancy and allowed to acclimate in the animal facility until the experiment was performed. In most cases and unless otherwise stated, experiments occurred on day 17 of a 21–22 day rat pregnancy.

Bacterial preparation

Bacteria (a pathogenic strain of *E. coli* originally isolated from a patient with urosepsis and obtained from the American Type Culture Collection (ATCC #12014)) were stored in 50% glycerol at -80°C and processed as previously described [4,23]. Live and heat-killed bacteria from this strain have been used repeatedly by our group to induce preterm delivery in mice.

For experiments with killed bacteria, a small chip of the stock culture was thawed in Luria-Bertani (LB) medium, passaged 4 times prior to overnight culture in 4 L of medium and concentrated by centrifugation. The bacteria were then washed three times in phosphate-buffered saline (PBS) and resuspended in 8 ml of PBS, yielding a concentration of 1.7×10^{11} bacteria per ml. This dense live bacterial prep was quantitated by plating serial dilutions for overnight culture just prior to the remaining stock being killed by boiling for 5 minutes. Non-viability of this boiled preparation was confirmed by overnight incubation of both plate and broth cultures. The stock of killed *E. coli* was aliquoted and frozen at -80°C. Prior to each experiment using killed bacteria, an aliquot was thawed and diluted in PBS to the desired concentration.

For experiments with live bacteria, *E. coli* were taken either from the frozen stock or from fresh plates and passaged twice, including one overnight culture in 3 ml of LB medium. At 10 AM, 100 µl of the overnight culture was added to 5 ml of sterile medium and grown at 37°C for 4

hours, when surgeries were performed. This bacterial prep was used undiluted and was kept at room temperature during surgery. The culture was quantified by plating before the first procedure and after the last procedure of the day, and the pre- and post-procedure counts were averaged to determine the inoculum used.

Surgical procedures

Rats were anesthetized with a mixture of 10 mg/kg Ketamine and 1 mg/kg Xylazine intraperitoneally. They were then immobilized in the supine position with the external genitalia slightly overhanging the edge of the lab bench. A 3 mm lacrimal duct scope attached to an external light source was advanced into the vagina. Polyethylene tubing (Becton Dickinson #427416, I.D. 0.76 mm (.030") O.D. 1.22 mm (.048")) was joined to the barrel of the scope through a narrow channel created from strips of tape. The tubing could be advanced or withdrawn for cannulation of the cervix under direct visualization. The cannula (primed with injectate to remove air) was advanced for a distance of approximately 1 cm through the cervical opening prior to injection of 100 – 1000 µl of solution. Various infusion volumes were used to assess their potential differential effects. After an average of 30 seconds (to allow for fluid distribution through the uterine horn while minimizing egress through the cervix), the cannula was withdrawn and visualization continued for another 30 seconds to assess whether fluid leakage occurred. The rat was then returned to her cage and allowed to awaken from anesthesia.

Original attempts using both an external pressurized air supply and a hand-operated bulb to distend the vagina to ease visualization of the cervix were abandoned due to air entry into the uterine horns. Thereafter, we were easily able to visualize and catheterize the cervical orifice without any source of pressurized air.

In general, animals tolerated these procedures well. Four rats died within 20 hours of surgery (presumably from anesthetic complications, as no specific findings were seen on necropsy). An additional 2 animals developed a severe shock-like syndrome, again with no findings on necropsy. These 6 subjects were excluded from the analysis.

In some cases, direct intrauterine injection of bacteria was performed at laparotomy to assess the effects of route of administration. These abdominal procedures were fashioned after our existing mouse model [4,23]. A 2 cm mid-line incision was made in the lower abdomen and the right uterine horn was exposed. Bacteria or sterile medium were then injected in volumes of 100 – 1000 µl into the midsection of the right uterine horn at a site between two adjacent fetuses. The peritoneal cavity was then closed

with interrupted sutures and the skin was closed with staples.

Surgical procedures lasted 10 – 15 minutes. Animals recovered in individual, clean cages in the animal facility and underwent twice-daily observations. The number of live-born or dead pups was noted. Preterm delivery was defined as the finding of at least one fetus in the cage or in the lower vagina within 48 hours of surgery.

In some cases, early euthanasia and necropsy were performed to assess catheter placement, dye distribution or fetal viability. Fetal viability was assessed by color, spontaneous movement and visible vascular pulsations in either the chest cavity or the vessels within fetal membranes. Other animals were observed for various periods of time through one week after delivery to record other observations, such as time to delivery and viability of delivered pups. Animals were euthanized by CO₂ inhalation plus thoracotomy.

Statistics

Statistical analysis of categorical values was by chi-square or contingency tables, with Fisher exact correction as needed.

Results

Development of a trans-cervical uterine infusion model in the rat

A total of 9 rats on days 14–18 of pregnancy underwent catheterization and immediate euthanasia to assess catheter position. A separate group of 14 pregnant rats on gestational days 17–19 (7 rats on day 17, 6 rats on day 18 and 1 rat on day 19) was instilled with PBS with or without sterile bromphenol blue dye in a total volume of 100–500 µl and observed for up to 70 hours to characterize the distribution of dye and any fetal effects.

After an initial learning period, we were able consistently to catheterize one uterine horn to the desired distance of 1 cm and infuse it with the dye solution with retention of dye in the uterine horn, continued viability of fetuses (verified by both immediate and delayed necropsy) and minimal leakage out the cervix and vagina. No control

animals delivered within 48 hours, the previously determined cutoff for preterm delivery (n = 14), and 11 of 12 rats followed to term delivered healthy litters (averaging 11.1 ± 2.7 pups per litter). The 12th rat had vaginal bleeding 20 hours after surgery and upon necropsy was found to have 6 absorbing conceptuses in one uterine horn along with 10 empty implantation sites distributed in both horns, suggestive of prior delivery from these sites. Therefore, we concluded that the trans-cervical infusion of various volumes of sterile solution was feasible and rarely affected the course of pregnancy.

Trans-cervical intrauterine infusion of a pathogenic strain of killed *E. coli* fails to induce preterm delivery

The trans-cervical infusion method was used to instill 2 × 10⁹ – 8 × 10¹⁰ killed *E. coli* organisms into the uteri of 14 pregnant rats at 17 days of gestation (Table 1). None of these animals delivered within 48 hours. Among these pregnancies, 6 were euthanized and necropsied at 48 hours to assess fetal status (see results below). The remaining 8 rats were observed for 70 hours without any preterm deliveries. No animals exhibited signs of systemic illness (such as piloerection, decreased mobility or decreased feces production).

Intrauterine infusion of live *E. coli* fails to induce preterm delivery

Our existing mouse model is characterized by a dramatic difference in potency of killed versus live bacteria for induction of preterm delivery (live bacteria are more potent by 5–6 orders of magnitude, with as few as 2000 organisms leading to preterm delivery) [4]. Therefore, we also experimented with live bacteria in an attempt to maximize the potential for inducing preterm delivery in rats. Eight pregnant rats were inoculated trans-cervically with 1 – 4 × 10⁸ live *E. coli* organisms. None of these animals delivered prematurely, including 3 animals that were followed to term, each of whom delivered live pups. None of the rats treated with live bacteria showed signs of systemic illness.

Finally, in order to investigate the effect of method of inoculation, 5 pregnant females were inoculated with 2 – 4 × 10⁸ live *E. coli* organisms by direct injection into the

Table 1: Effect of intrauterine *E. coli* inoculation on preterm delivery.

Treatment	Number of <i>E. coli</i> organisms	Number of animals	Preterm delivery rate (%)
Trans-cervical infusion			
Control	N/A	16	0
Killed <i>E. coli</i>	0.2 – 8 × 10 ¹⁰	14	0
Live <i>E. coli</i>	1 – 4 × 10 ⁸	8	0
Infusion at laparotomy			
Control	N/A	1	0
Live <i>E. coli</i>	2–7 × 10 ⁸	5	0

'Preterm' was defined as 48 hours after surgery, however some animals were observed for longer periods, up to term (see text).

right uterine horn at laparotomy. None of these 5 rats delivered prematurely.

Fetal effects of intrauterine bacterial infusion

The effect of intrauterine bacterial infusion on fetal viability was assessed by timed necropsy after surgery. One hundred percent of 80 fetuses from 6 mothers who received control (non-bacterial) injections and were euthanized 24 – 70 hours later were still alive at the time of necropsy. Also, as noted above, dams followed to term after control injections delivered large, healthy litters. In 14 litters treated with $2 \times 10^9 - 8 \times 10^{10}$ killed bacteria, 93% of 182 fetuses were alive at necropsy 48 – 70 hours later ($p = 0.014$, comparing control to killed bacterial injection). There were no significant trends associating inoculum size with fetal survival. Thus, killed *E. coli* has a statistically significant though small impact on fetal viability.

In contrast, live bacteria had a large effect on fetal viability that was dependent upon route of administration. Trans-cervical inoculation with live *E. coli* was associated with

survival at necropsy 48 hours later in 64% of 14 fetuses, 47% of 28 fetuses and 32% of 31 fetuses from 5 pregnant rats administered 7×10^7 , 2×10^8 or 4×10^8 live *E. coli*, respectively ($p = 0.13$) (Figure 1). Direct administration of live bacteria by intrauterine injection at laparotomy produced even more profound effects on fetal survival, with only 3.4% of 63 fetuses from 5 pregnant dams still alive at necropsy 48 hours later ($p = 0.0006$ for comparison of the trans-cervical and direct injection routes of administration of 2×10^8 bacteria and $p = 0.003$ for comparison of the two routes of administration of 4×10^8 bacteria). Finally, each of 3 pregnant dams who received $1 - 2 \times 10^8$ live *E. coli* trans-cervically and were followed through spontaneous delivery produced live pups at term, but their number was diminished (average 3 per litter), compared with 10.2 pups per litter in control animals ($p = 0.001$).

Discussion

The present study describes the inability to induce pre-term delivery in pregnant rats using live or killed *E. coli*. This experience stands in sharp contrast to published




Figure 1

Effect of intrauterine treatment on fetal viability. Female rats on day 17 of pregnancy were inoculated with either sterile solution (control), killed *E. coli* or live *E. coli* bacteria, by either trans-cervical infusion or direct intrauterine injection at laparotomy. Animals were euthanized 24–70 hours later and fetal viability was determined. Displayed on the Y-axis is the percentage of fetuses alive at the time of autopsy. The number of treated dams in each group is also displayed.

experience with mice, in which live or killed *E. coli* as well as multiple other inflammatory/infectious stimuli produced preterm delivery [3,4,6-9,24,25]. These stimuli include a variety of bacteria, pathogen extracts (e.g. LPS, peptidoglycan, lipoteichoic acid, poly inosinic:cytidylic acid), cytokines (e.g. IL-1, TNF), and others. The strain of *E. coli* used in the present study has been used successfully in models of preterm birth not only by our group but by others also in mice [8,26] and rabbits [12,27].

We could identify in the literature only two laboratories that reported induction of preterm delivery using an infectious/inflammatory stimulus in rats [20-22]. In work from the first group, Sprague Dawley rats underwent intrauterine implantation of catheters on day 15 or 16 followed on day 17 of pregnancy by infusion of 25 µg or 50 µg of LPS over 4 hours. Saline treatment was followed by delivery in an average of 117 hours, whereas 25 µg and 50 µg of LPS produced delivery in 82 and 63 hours, respectively, together with increased production of prostaglandin F_{2α} metabolite in placentas of LPS-exposed animals [20]. In a separate study from the same group, a 50 µg infusion of LPS produced preterm delivery in an average of 92 hours, which was approximately one day before delivery in saline-treated rats [22]. Treatment with IL-10 restored delivery to term. It is not clear why there was a discrepancy of nearly 30 hours in the interval to delivery between these two studies in rats treated with 50 µg LPS. In a comment, the authors note that slow infusion over 4 hours appears to be important for 'optimal results' [20]. For our studies we chose 48 hours as a reasonable cut-off for 'preterm delivery' to avoid approaching term and because this cutoff is practical in the mouse (in which delivery in similar models usually occurs in less than 24 hours). When delivery did not occur in this time frame, we extended our observations to term and still did not observe early delivery.

In work from the second group, Wistar rats were treated with 25 µg/kg of *E. coli* LPS intraperitoneally on day 16 of pregnancy [21]. Although no animals in this study were treated with control injections, the average time to delivery was approximately 22 hours. It should be noted that assuming an average weight of 333 gm per rat, these subjects would have received 8.3 µg of LPS each. Somewhat surprisingly, intraperitoneal treatment with erythromycin significantly prolonged the interval to delivery, by up to 50 hours.

Whether the different potencies reflected in the above two experimental designs is due to route of administration (intrauterine versus intraperitoneal), rate of infusion, rat strain (Sprague-Dawley versus Wistar) or some other factor is not clear. In the present report, Sprague Dawley rats treated with $2 \times 10^9 - 8 \times 10^{10}$ killed *E. coli* would have

received between 20 µg and 800 µg of LPS each, assuming that each *E. coli* bacterial cell contains 9.7×10^{-15} gm LPS [28].

Far more numerous than the 3 studies cited above that demonstrate preterm birth after LPS administration are the studies that use hormonal manipulation (primarily progesterone antagonism) to induce preterm delivery in rats (examples to be found in [18,29,30]). This notable fact is reinforced by another, namely that the literature contains several examples of experimental LPS exposure in rats after which observation periods of sufficient length were conducted to effectively rule out subsequent preterm delivery. For example, *E. coli*-derived LPS (400 µg/kg intraperitoneally) induced cervical changes in Sprague Dawley rats when administered on day 14 of pregnancy [30]. Animals were harvested 2 days after treatment with no cases of preterm delivery reported. In a study of the effect of prenatal LPS on neurobehavioral development, pregnant Wistar rats were administered 2 mg/kg of *E. coli* LPS subcutaneously on each day of pregnancy from conception through delivery [31]. Though maternal LPS treatment induced significant changes in neonatal neurobehavioral outcomes as well as dopaminergic neurotransmission and synaptophysin expression, there was no reported effect on timing of delivery. A different study in which Sprague Dawley rats were treated with LPS (100 µg/kg intraperitoneally) every day from day 14 to day 20 of pregnancy demonstrated a significant increase in fetal death and reduced size of surviving fetuses compared with controls [32]. Within the uteri of treated dams there was increased apoptosis, a doubling of TNFα levels and a tripling of nitric oxide and myeloperoxidase levels, however no preterm labor was reported. In a study focusing on fetal/neonatal brain injury, *E. coli* LPS treatment (500 µg/kg intraperitoneally) of pregnant Sprague Dawley rats on day 18 and again on day 19 of pregnancy produced white matter injury, myelination defects and apoptosis and increased expression of IL-1β, TNFα and IL-6 in pup brains 7 days after birth [33]. Again, no preterm delivery was reported. In a Chinese language publication available in English in abstract form, treatment of Sprague Dawley rats with intraamniotic LPS (quantity not provided) on day 15 of pregnancy produced both histologic changes and reduction in expression of vascular endothelial growth factor (VEGF) and its receptors (Flk-1 and Flt-1) in neonatal lungs, but animals delivered at term [34].

Why has the induction of inflammation-induced preterm birth been so difficult and variable in the rat, while it is so readily accomplished in mice and other species? The results cited above and the fact that in the present study there were measurable effects on fetal survival suggest that the pregnant Sprague Dawley strain of rat is responsive to *E. coli*. Among the demonstrable effects of LPS in pregnant

rats are uterine apoptosis and increased expression of TNF α , nitric oxide and myeloperoxidase [32], increased prostaglandins in the amniotic fluid and an increase in *in vitro* myometrial contractile activity in response to oxytocin [35], and an increase in both spontaneous and agonist-induced uterine contractions in myometrial strips isolated from LPS-treated rats [36]. Therefore, it seems likely that failure to induce preterm delivery in this model of bacterial infection is specific for parturition, rather than due to a global hyporesponsiveness to either Gram negative bacteria or LPS.

Among the possible explanations for this phenomenon is that LPS may have a differential effect among different rodent species with respect to parturition-related factors. A number of genes have been identified in the mouse that are critical for a normal response in parturition models. It is possible that some of these genes or others not yet identified are differentially expressed in various rat strains, thus accounting for a lack of responsiveness. Further comparative analysis of genomic sequence information (particularly between mice and rats) may provide valuable insights in this regard. Terrone noted that high doses of LPS induced intrauterine fetal demise, while lower doses tended to produce fewer fetal deaths and preterm delivery [22]. Two groups have reported that LPS increases uterine myometrial contractility in rat uterus in response to oxytocin [35,36]. However, spontaneous uterine contractions induced by LPS in the rat do not appear to be mediated by prostaglandins, as prostaglandin inhibitors fail to diminish this response [36]. This appears contrary to relevant mechanisms in other species. Finally, uterine contractions in the rat may be dissociated from cervical ripening. In one study, a platelet-activating factor (PAF-) antagonist blocked cervical changes induced by LPS but not those induced by the antiprogesterin RU-486 [30]. Perhaps the rat laboring after exposure to LPS or *E. coli*, unlike the spontaneously laboring rat at term, does not experience the kind of cervical modifications ('ripening') that allow it to deliver.

We observed a difference in the fetal toxicity of a live bacterial inoculum between trans-cervical inoculation with a catheter and direct intrauterine inoculation at laparotomy. We suspect that this difference may be due to delayed leakage of the trans-cervical inoculum through the dilated cervical orifice left behind by the catheterization procedure, thus effectively diminishing the size of the inoculum.

It is our hope that publication of the present negative results will either spur successful efforts to create an improved rat model or lead to potentially useful insights as to why this rodent behaves so differently from the mouse. Such insights have the potential to lead to appli-

cations in humans that will diminish the burden of pre-mature birth.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

EH contributed to the conception and design, development of the rat model, analysis of data, and drafting of the manuscript. YF contributed to the development of the rat model, carried out the experiments, analyzed results and helped draft the manuscript. RR contributed to the conception and design, analysis of data, and drafting of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Mr. Lennard Wahlberg provided technical assistance with development of the trans-cervical infusion model.

This research was supported in part by the Intramural Research Program of the National Institute of Child Health and Human Development, NIH, DHHS.

References

- Klein LL, Gibbs RS: **Use of microbial cultures and antibiotics in the prevention of infection-associated preterm birth.** *Am J Obstet Gynecol* 2004, **190(6)**:1493-1502.
- Elovitz MA, Mrinalini C: **Animal models of preterm birth.** *Trends Endocrinol Metab* 2004, **15(10)**:479-487.
- Fidel PL, Romero R, Wolf N, Cutright J, Ramirez M, Arana H, Cotton DB: **Systemic and local cytokine profiles in endotoxin-induced preterm parturition in mice.** *Am J Obstet Gynecol* 1994, **170**:1467-1475.
- Hirsch E, Saotome I, Hirsh D: **A model of intrauterine infection and preterm delivery in mice.** *Am J Obstet Gynecol* 1995, **172(5)**:1598-1603.
- Hirsch E, Wang H: **The molecular pathophysiology of bacterially induced preterm labor: insights from the murine model.** *J Soc Gynecol Investig* 2005, **12(3)**:145-155.
- Ilievski V, Lu SJ, Hirsch E: **Activation of toll-like receptors 2 or 3 and preterm delivery in the mouse.** *Reprod Sci* 2007, **14(4)**:315-320.
- Kajikawa S, Kaga N, Futamura Y, Kakinuma C, Shibutani Y: **Lipoteichoic acid induces preterm delivery in mice.** *J Pharmacol Toxicol Methods* 1998, **39(3)**:147-154.
- Reznikov LL, Fantuzzi G, Selzman CH, Shames BD, Barton HA, Bell H, McGregor JA, Dinarello CA: **Utilization of endoscopic inoculation in a mouse model of intrauterine infection-induced preterm birth: role of interleukin 1beta.** *Biol Reprod* 1999, **60(5)**:1231-1238.
- Romero R, Mazor M, Tartakovsky B: **Systemic administration of interleukin-1 induces preterm parturition in mice.** *Am J Obstet Gynecol* 1991, **165(4)**:969-971.
- Bry K, Hallman M: **Transforming growth factor- β 2 prevents preterm delivery induced by interleukin-1 α and tumor necrosis factor- α in the rabbit.** *Am J Obstet Gynecol* 1993, **168(4)**:1318-1322.
- Davies JK, Shikes RH, Sze CI, Leslie KK, McDuffie RS Jr, Romero R, Gibbs RS: **Histologic inflammation in the maternal and fetal compartments in a rabbit model of acute intra-amniotic infection.** *Am J Obstet Gynecol* 2000, **183(5)**:1088-1093.
- Dombroski RA, Woodard DS, Harper MJ, Gibbs RS: **A rabbit model for bacteria-induced preterm pregnancy loss.** *Am J Obstet Gynecol* 1990, **163(6)**:1938-1943.
- Gibbs RS, McDuffie RS Jr, Kunze M, Barr JM, Wolf DM, Sze CI, Shikes R, Sherman MP: **Experimental intrauterine infection with**

- Prevotella bivia** in New Zealand White rabbits. *Am J Obstet Gynecol* 2004, **190(4)**:1082-1086.
14. Gravett MG, Adams KM, Sadowsky DW, Grosvenor AR, Witkin SS, Axthelm MK, Novy MJ: **Immunomodulators plus antibiotics delay preterm delivery after experimental intraamniotic infection in a nonhuman primate model.** *Am J Obstet Gynecol* 2007, **197(5)**:518 e511-518.
 15. Gravett MG, Witkin SS, Haluska GJ, Edwards JL, Cook MJ, Novy MJ: **An experimental model for intraamniotic infection and preterm labor in rhesus monkeys.** *Am J Obstet Gynecol* 1994, **171(6)**:1660-1667.
 16. Sadowsky DW, Adams KM, Gravett MG, Witkin SS, Novy MJ: **Preterm labor is induced by intraamniotic infusions of interleukin-1beta and tumor necrosis factor-alpha but not by interleukin-6 or interleukin-8 in a nonhuman primate model.** *Am J Obstet Gynecol* 2006, **195(6)**:1578-1589.
 17. Buhimschi C, Boyle MB, Saade GR, Garfield RE: **Uterine activity during pregnancy and labor assessed by simultaneous recordings from the myometrium and abdominal surface in the rat.** *Am J Obstet Gynecol* 1998, **178(4)**:811-822.
 18. Garfield RE, Gasc JM, Baulieu EE: **Effects of the antiprogesterone RU 486 on preterm birth in the rat.** *Am J Obstet Gynecol* 1987, **157(5)**:1281-1285.
 19. Yallampalli C, Buhimschi I, Chwalisz K, Garfield RE, Dong YL: **Preterm birth in rats produced by the synergistic action of a nitric oxide inhibitor (NG-nitro-L-arginine methyl ester) and an antiprogesterone (onapristone).** *Am J Obstet Gynecol* 1996, **175(1)**:207-212.
 20. Bennett WA, Terrone DA, Rinehart BK, Kassab S, Martin JN Jr, Granger JP: **Intrauterine endotoxin infusion in rat pregnancy induces preterm delivery and increases placental prostaglandin F2alpha metabolite levels.** *Am J Obstet Gynecol* 2000, **182(6)**:1496-1501.
 21. Celik H, Ayar A: **Effects of erythromycin on pregnancy duration and birth weight in lipopolysaccharide-induced preterm labor in pregnant rats.** *Eur J Obstet Gynecol Reprod Biol* 2002, **103(1)**:22-25.
 22. Terrone DA, Rinehart BK, Granger JP, Barrilleaux PS, Martin JN Jr, Bennett WA: **Interleukin-10 administration and bacterial endotoxin-induced preterm birth in a rat model.** *Obstet Gynecol* 2001, **98(3)**:476-480.
 23. Mussalli GM, Blanchard R, Brunnert SR, Hirsch E: **Inflammatory cytokines in a murine model of infection-induced preterm labor: cause or effect?** *J Soc Gynecol Invest* 1999, **6**:188-195.
 24. Fidel PJ Jr, Romero R, Maymon E, Hertelendy F: **Bacteria-induced or bacterial product-induced preterm parturition in mice and rabbits is preceded by a significant fall in serum progesterone concentrations.** *J Matern Fetal Med* 1998, **7(5)**:222-226.
 25. Robertson SA, Skinner RJ, Care AS: **Essential role for IL-10 in resistance to lipopolysaccharide-induced preterm labor in mice.** *J Immunol* 2006, **177(7)**:4888-4896.
 26. Haddad R, Gould BR, Romero R, Tromp G, Farookhi R, Edwin SS, Kim MR, Zingg HH: **Uterine transcriptomes of bacteria-induced and ovariectomy-induced preterm labor in mice are characterized by differential expression of arachidonate metabolism genes.** *Am J Obstet Gynecol* 2006, **195(3)**:822-828.
 27. Hasegawa A, Otsuki K, Sasaki Y, Sawada M, Mitsukawa K, Chiba H, Nagatsuka M, Okai T, Kato A: **Preventive effect of recombinant human lactoferrin in a rabbit preterm delivery model.** *Am J Obstet Gynecol* 2005, **192(4)**:1038-1043.
 28. Neidhardt FC: **Chemical Composition of Escherichia coli.** In *Escherichia coli and Salmonella typhimurium: cellular and molecular biology Volume 1*. Edited by: Neidhardt FC. Washington, DC: American Society for Microbiology; 1987:1654.
 29. Cirillo R, Tos EG, Page P, Missotten M, Quattropiani A, Scheer A, Schwarz MK, Chollet A: **Arrest of preterm labor in rat and mouse by an oral and selective nonprostanoid antagonist of the prostaglandin F2alpha receptor (FP).** *Am J Obstet Gynecol* 2007, **197(1)**:54 e51-59.
 30. Maul H, Shi L, Marx SG, Garfield RE, Saade GR: **Platelet-activating factor antagonist WEB-2170 inhibits lipopolysaccharide-induced, but not antiprogesterone-induced, preterm cervical ripening in timed-pregnant rats.** *Am J Obstet Gynecol* 2003, **189(4)**:963-967.
 31. Romero E, Ali C, Molina-Holgado E, Castellano B, Guaza C, Borrell J: **Neurobehavioral and immunological consequences of pre-natal immune activation in rats. Influence of antipsychotics.** *Neuropsychopharmacology* 2007, **32(8)**:1791-1804.
 32. Rivera DL, Ollister SM, Liu X, Thompson JH, Zhang XJ, Pennline K, Azuero R, Clark DA, Miller MJ: **Interleukin-10 attenuates experimental fetal growth restriction and demise.** *Faseb J* 1998, **12(2)**:189-197.
 33. Kumral A, Baskin H, Yesilirmak DC, Ergur BU, Aykan S, Genc S, Genc K, Yilmaz O, Tugyan K, Giray O, et al.: **Erythropoietin attenuates lipopolysaccharide-induced white matter injury in the neonatal rat brain.** *Neonatology* 2007, **92(4)**:269-278.
 34. Wang W, Wei W, Ning Q, Luo XP: **[Effect of intra-amniotic endotoxin priming plus hyperoxic exposure on the expression of vascular endothelial growth factor and its receptors in lungs of preterm newborn rats].** *Zhonghua Er Ke Za Zhi* 2007, **45(7)**:533-538.
 35. Okawa T, Suzuki H, Yaanagida K, Sato A, Vedernikov Y, Saade G, Garfield R: **Effect of lipopolysaccharide on uterine contractions and prostaglandin production in pregnant rats.** *Am J Obstet Gynecol* 2001, **184(2)**:84-89.
 36. Ross RG, Sathishkumar K, Naik AK, Bawankule DU, Sarkar SN, Mishra SK, Prakash VR: **Mechanisms of lipopolysaccharide-induced changes in effects of contractile agonists on pregnant rat myometrium.** *Am J Obstet Gynecol* 2004, **190(2)**:532-540.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

