

5-1-2011

Bayesian Threshold Moving Average Models


Mahmoud M. Smadi

Jordan University of Science and Technology, smadi@just.edu.jo

M. T. Alodat

Yarmouk University, Irbid, Jordan, alodatmts@yahoo.com

Follow this and additional works at: <http://digitalcommons.wayne.edu/jmasm>

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

Recommended Citation

Smadi, Mahmoud M. and Alodat, M. T. (2011) "Bayesian Threshold Moving Average Models," *Journal of Modern Applied Statistical Methods*: Vol. 10 : Iss. 1 , Article 23.

DOI: 10.22237/jmasm/1304223720

Available at: <http://digitalcommons.wayne.edu/jmasm/vol10/iss1/23>

This Regular Article is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Journal of Modern Applied Statistical Methods by an authorized editor of DigitalCommons@WayneState.

Bayesian Threshold Moving Average Models

Mahmoud M. Smadi
Jordan University of Science and Technology,
Irbid, Jordan

M. T. Alodat
Yarmouk University,
Irbid, Jordan

A Bayesian approach in threshold moving average model for time series with two regimes is provided. The posterior distribution of the delay and threshold parameters are used to examine and investigate the intrinsic characteristics of this nonlinear time series model. The proposed approach is applied to both simulated data and a real data set obtained from a chemical system.

Key words: Threshold time series, moving average model, Bayesian estimation, simulation, chemical data.

Introduction

One class of nonlinear time series models is the threshold time series models which are extensively reported in literature. Among these, Tong and Lim (1980) introduced threshold autoregressive (TAR) models with statistical inference and applications. Bayesian inference for threshold autoregressive models have been investigated by different authors.

Geweke and Terui (1991) derived an exact posterior distribution of the delay and threshold parameters. Cathy, et al. (1995) used Monte Carlo Markov chain (MCMC) methods to implement a Bayesian inference on TAR models, and Broemeling and Cook (1992) performed a Bayesian analysis on TAR models. However, most of the literature emphasizes the threshold autoregressive models. Wang, et al. (1984) introduced the threshold autoregressive

moving average (TARMA) models and considered the estimation of the model parameters. De Gooijer (1998) studied various problems associated with the identification, estimation and testing of threshold moving average models. Ling and Tong (2005) considered a quasi-likelihood ratio test for the threshold in moving average models. Amendola, et al. (2009) discussed the stochastic structure of the self-exiting TARMA model; they specified sufficient conditions for weak stationarity and showed that the self-exiting TARMA model belongs to the class of the random coefficients autoregressive models. Smadi (1997) used the Bayesian approach for exploration of the joint posterior distribution for TARMA models using MCMC methods: he assumed noninformative priors, fixing the delay parameter d . In addition, he used a modified Gibbs sampling scheme, which is a hybrid strategy of Gibbs sampler, random walk Metropolis, and importance sampling. Safadi and Morettin (2000) considered a Bayesian analysis for threshold autoregressive moving average models and a hierarchical prior to perform Bayesian analysis using a rearranged procedure with MCMC methods.

The objective of this study is to provide a Bayesian approach in a threshold moving average model for time series with two regimes. The posterior distribution of the delay and the threshold parameters are used to examine and investigate the characteristics which are intrinsic to this nonlinear time series model. The

Mahmoud M. Smadi is an Associate Professor of Statistics in the Department of Mathematics and Statistics. His research interests include time series analysis, Bayesian statistics, and environmental statistics. Email him at: smadi@just.edu.jo. M. T. Alodat is an Associate Professor in the Department of Statistics. His research interests include the geometry of random fields, analysis medical images, approximating probability distributions, prediction and estimation using Bayesian methods. Email him at: alodatmts@yahoo.com.

proposed approach is applied to both simulated data and a real data set obtained from a chemical system.

Methodology

The Threshold Models

Let $\{Y_t, t \geq 1\}$ be a time series, the threshold autoregressive moving average models with two regimes. Wang, et al. (1984) symbolized $TARMA(2, (p_1, q_1), (p_2, q_2))$, given by:

$$Y_t = \begin{cases} \sum_{i=1}^{p_1} \phi_i^{(1)} Y_{t-i} + \sum_{i=1}^{q_1} \theta_i^{(1)} Z_{t-i}^{(1)} + Z_t^{(1)}, & Y_{t-d} \leq r, \\ \sum_{i=1}^{p_2} \phi_i^{(2)} Y_{t-i} + \sum_{i=1}^{q_2} \theta_i^{(2)} Z_{t-i}^{(2)} + Z_t^{(2)}, & Y_{t-d} > r, \end{cases} \quad (1)$$

where $\phi_1^{(j)}, \dots, \phi_{p_j}^{(j)}, \theta_1^{(j)}, \dots, \theta_{q_j}^{(j)}, (j=1,2)$ are the model coefficients of the two ARMA subsystems, r is called the threshold parameter and d is the delay parameter; assume the innovations $Z_t^{(j)} : N(0, \sigma_j^2), j=1, 2$.

A special case of equation (1), is the threshold autoregressive model $TAR(2; p_1, p_2)$:

$$Y_t = \begin{cases} \sum_{i=1}^{p_1} \phi_i^{(1)} Y_{t-i} + Z_t^{(1)} & Y_{t-d} \leq r, \\ \sum_{i=1}^{p_2} \phi_i^{(2)} Y_{t-i} + Z_t^{(2)} & Y_{t-d} > r. \end{cases} \quad (2)$$

Another special case of equation (1) is the threshold Moving Average model $TMA(2; q_1, q_2)$:

$$Y_t = \begin{cases} \sum_{i=1}^{q_1} \theta_i^{(1)} Z_{t-i}^{(1)} + Z_t^{(1)} & Y_{t-d} \leq r, \\ \sum_{i=1}^{q_2} \theta_i^{(2)} Z_{t-i}^{(2)} + Z_t^{(2)} & Y_{t-d} > r. \end{cases} \quad (3)$$

Posteriors Distribution

The approximate posterior distribution of the delay and threshold parameters (d, r) for threshold moving average models (3) is based on using estimated residuals instead of the true

innovations. Broemeling and Shaarawy (1986) implemented the estimated innovations for Bayesian analysis of ARMA models. Smadi (1997) and Safadi and Morettin (2000) have used this estimated innovation approach to explore the posterior distributions of the threshold autoregressive moving average models.

Defining $\phi = (d, r)$, and the set $B_j = (\theta_1^{(j)}, \theta_2^{(j)}, \dots, \theta_{q_j}^{(j)})$, and under the normality assumption, that is, $Z_t^{(j)} i.i.d N(0, \sigma_j^2)$, a prior $\pi(\phi)$ could be any form as long as

$$\sum_{d=1}^R \int_{-\infty}^{\infty} \pi(d, r) dr = 1.$$

Conditional on ϕ , independent priors on B_j and σ_j of standard Jeffreys prior can be expressed as:

$$\prod \{B_j, \sigma_j (j=1,2), \phi\} = \pi_c(B_j, \sigma_j (j=1,2), \phi) \pi(\phi) \quad (4)$$

where $\pi_c(B_j, \sigma_j, j=1,2 | \phi) = (\sigma_1 \sigma_2)^{-1}$. It is also assumed that $Z_0^{(1)} = Z_{-1}^{(1)} = \dots = Z_{1-q_1}^{(1)} = 0$ and $Z_0^{(2)} = Z_{-1}^{(2)} = \dots = Z_{1-q_2}^{(2)} = 0$. Conditioning on $\phi = (d, r)$, estimates of the innovations $Z_t^{(1)}$ and $Z_t^{(2)}$ can be obtained using least squares estimates. In this case, the following model is obtained:

$$Y_t = \begin{cases} \sum_{i=1}^{q_1} \theta_i^{(1)} \widehat{Z}_{t-i}^{(1)} + Z_t^{(1)} & Y_{t-d} \leq r, \\ \sum_{i=1}^{q_2} \theta_i^{(2)} \widehat{Z}_{t-i}^{(2)} + Z_t^{(2)} & Y_{t-d} > r. \end{cases} \quad (5)$$

Derivation of the approximate posterior of the delay and the threshold parameters of the threshold moving average model is similar to the threshold autoregressive model (2) reported by Geweke and Terui (1993). After estimating the

innovations and using the model (3), conditional on $\phi = (d, r)$, let W_1 be a vector consisting of N_1 ordered observations on $\{Y_t\}$ such that $Y_{t-d} \leq r$, and let W_2 be a vector consisting of N_2 ordered observations on $\{Y_t\}$ such that $Y_{t-d} > r$. Let X_j be a $N_j \times q_j$ matrix of lagged variables on the estimated innovations $\{\hat{Z}_t\}$ corresponding to W_j ($j = 1, 2$). Then, the approximate posterior density of $(B_1, \sigma_1, B_2, \sigma_2)$ conditional on d and r is the product of two posterior densities, that is:

$$P(B_j, \sigma_j, j = 1, 2 | \phi, Y) = \prod_{j=1}^2 \sigma_j^{-(N_j+1)} (2\pi)^{-(N_j/2)} \exp \left\{ -\frac{(W_j - X_j B_j)' (W_j - X_j B_j)}{2\sigma_j^2} \right\} \quad (6)$$

The posterior distribution of ϕ can be derived by integrating this expression with respect to $(B_1, \sigma_1, B_2, \sigma_2)$. The problem is to integrate the following expression with respect to σ :

$$\sigma^{-(T+1)} \exp \left\{ -\frac{(W - XB)' (W - XB)}{2\sigma^2} \right\} = \sigma^{-(T+1)} \exp \left\{ -\frac{v s^2 + (B - \hat{B})' X' X (B - \hat{B})}{2\sigma^2} \right\} \quad (7)$$

where

$$B = (X' X)^{-1} X' W, \quad s^2 = (W - X\hat{B})' (W - X\hat{B}) / v, \quad v = N - K - 1.$$

Integrating over σ_1 and σ_2 on the right hand side of equation (7), it is possible to obtain:

$$P\{B_j, j = 1, 2 | \phi, Y\} = \prod_{j=1}^2 (2\pi)^{-(N_j/2)} \int_0^\infty \sigma_j^{-(N_j+1)} \exp \left\{ -\frac{v_j s_j^2 + (B_j - \hat{B}_j)' W_j' W_j (B_j - \hat{B}_j)}{2\sigma_j^2} \right\} d\sigma_j$$

$$P\{B_j, j = 1, 2 | \phi, Y\} = \prod_{j=1}^2 (2\pi)^{-(N_j/2)} 2^{(N_j/2-1)} \Gamma\left(\frac{N_j}{2}\right) \{v_j s_j^2 + (B_j - \hat{B}_j)' W_j' W_j (B_j - \hat{B}_j)\}^{-(N_j/2)} \quad (8)$$

multiplying equation (8) by $\pi(\phi)$ and integrating B_j , for $j = 1, 2$ out results in the posterior distribution of ϕ (Geweke & Terui, 1993):

$$P(\phi | Y) \propto \pi(\phi) \prod_{j=1}^2 (2\pi)^{-(N_j/2)} 2^{(N_j/2-1)} \Gamma\left(\frac{N_j}{2}\right) (s_j^2)^{-(N_j/2)} \times \int_{R^{(k_j+1)}} \left\{ v_j + (B_j - \hat{B}_j)' \frac{W_j' W_j}{s_j^2} (B_j - \hat{B}_j) \right\}^{-(N_j/2)} dB_j = \prod_{j=1}^2 2^{-(v_j/2+1)} (\pi)^{-(v_j/2)} \Gamma\left(\frac{v_j}{2}\right) \left(\frac{v_j s_j^2}{2}\right)^{-(v_j/2)} |W_j' W_j|^{-1/2} \quad (9)$$

where

$$s_j^2 = (W_j - X_j \hat{B}_j)' (W_j - X_j \hat{B}_j) / v_j, \quad \hat{B}_j = (X_j' X_j)^{-1} X_j' W_j, \quad v_j = N_j - K_j - 1.$$

Results

The characteristics of the posterior distribution of $\phi = (d, r)$ were investigated for simulated data and a real data set obtained from a chemical system. As a set of possible values of the threshold parameter r , the order statistics $[Y^{(i)}, i = 1, \dots, q(\leq N)]$ of observations was used.

Simulation Examples

Simulation results were based on both one realization and 100 realizations. The TMA(2;1,1) model was considered, where $\theta^{(1)} = -0.4, \theta^{(2)} = 0.4, \sigma_1^2 = 1, \sigma_2^2 = 1, d = 1$, and $r = 0$. A one realization was generated with series length of 50. As a set of possible values of the threshold parameter $r, [r_0, r_L]$ was chosen as large as possible and the delay parameter d was selected as $d = 1, 2, 3, 4$ and 5 .

Simulation results demonstrated that the posterior mass was concentrated at $d=1, 2$ and 3 . Summary results of the joint posterior distribution of $\phi = (d, r)$ are presented in Table 1. The marginal posterior distribution of $d = 1, 2, 3$ have probabilities of 0.5566, 0.2500 and 0.1800 respectively. The posterior probability concentrates predominantly on few points, namely $(d, r) = (1, 0.0036)$ and $(1, 0.0350)$ with respective probabilities of 0.05582 and 0.06987.

Simulation results based on 100 realizations with series length of 100 were analyzed. For each realization, $\phi = (d, r)$ was estimated based on modal value of the posterior distribution. The results yield relative frequencies of 87%, 7%, and 6% for $d = 1, 2,$ and 3 respectively. The marginal posterior of r is shown in Figure 1; as expected, the model value is concentrated around the true threshold value $r = 0$.

Real Data Example

Series A, which consists of 197 observations and represents the concentration of a chemical process, was considered (Box & Jenkins, 1976). The differenced time series was considered. Fitting MA(1) model yields

$$X_t = -0.69Z_{t-1} + Z_t \quad (15)$$

Smadi (1997) used the MCMC technique for exploration of the posterior distribution of the threshold parameter r . The methodology proposed herein is applied to the differenced series in order to examine the posterior distribution of the threshold and the delay parameter $\phi = (d, r)$. The number of threshold points is reduced from 196 to 22 points because some differences have the same values. Values of $[-0.4, 0.4]$ were assigned for $[r_0, r_L]$. For the delay parameter d , the set $d = 1, 2, 3, 4$ and 5 were selected. It was found that the posterior mass was concentrated at $d = 1, 2$ and 3 .

Summary results of the joint posterior distribution of $\phi = (d, r)$ are presented in Table 2. It can be seen that the marginal posterior distribution of $d = 1, 2, 3$ have probabilities of 0.8344, 0.13163 and 0.03398, respectively. Also, the posterior probability concentrates on

$(d, r) = (1, 0.0)$ with probability of 0.17; this corresponds to the largest mode of the posterior density. Conditioning on $(d, r) = (1, 0.0)$, the fitted TMA(2;1,1) model is

$$X_t = \begin{cases} -0.72Z_{t-1}^{(1)} + Z_t^{(1)} & \text{if } X_{t-1} \leq 0, \\ -0.66Z_{t-1}^{(2)} + Z_t^{(2)} & \text{if } X_{t-1} > 0. \end{cases}$$

where $\hat{\sigma}_1^2 = 0.15$ and $\hat{\sigma}_2^2 = 0.12$.

Conclusion

From the proposed methodology and numerical results it can be concluded that the threshold moving average models are tractable from a Bayesian point of view. The nonlinearity threshold-type for moving average models can be detected by examining the marginal posterior distribution of the threshold parameter.

References

- Amendola, A., Niglio, M., & Vitale C. (2009). Statistical properties of threshold models. *Communication in Statistics-Theory and Methods*, 38, 2479-2497.
- Box, G. E. P., & Jenkins, G. M. (1976). *Time Series Analysis, Forecasting and Control*. San Francisco, CA: Holden-Day.
- Shaarawy, S., & Bromeling, L. (1984). Bayesian inferences and forecasts with moving average processes. *Communication in Statistics: Theory and Methods*, 13(15), 1871-1888.
- Cathy, W. S., Chen, ,& Lee, J. C. (1995). Bayesian inference of threshold models, *Journal of Time Series Analysis*, 16(5), 483-491.
- Geweke, J., & Terui, N. (1993). Bayesian threshold autoregressive models for nonlinear time series. *Journal of Time Series Analysis*, 14(5), 441-54.
- Gooijer, J. D. (1998). On threshold moving-average models. *Journal of time series analysis*, 19(1), 1-18.
- Ling, S., & Tong, H. (2005). Testing for a linear MA model against threshold MA models. *Annals of Statistics*, 3, 2529-52.
- Safadi, T., & Morettin, P. A. (2000). Bayesian Analysis of threshold autoregressive moving average models. *Sankhya*, 62, Series B(3), 353-371.

BAYESIAN THRESHOLD MOVING AVERAGE MODELS

Table 1: Summary of Joint Posterior Densities of (r, d)

(r, d)	Integrated Density	(r, d)	Integrated Density	(r, d)	Integrated Density
(-1.1832, 1)	0.02234	(-1.1832, 2)	0.00345	(-1.1832, 3)	0.01233
(-1.0101, 1)	0.02278	(-1.0101, 2)	0.00263	(-1.0101, 3)	0.00771
(-0.8562, 1)	0.02282	(-0.8562, 2)	0.00223	(-0.8562, 3)	0.00513
(-0.7346, 1)	0.01532	(-0.7346, 2)	0.00192	(-0.7346, 3)	0.00510
(-0.6573, 1)	0.01058	(-0.6573, 2)	0.00181	(-0.6573, 3)	0.00321
(-0.6326, 1)	0.00839	(-0.6326, 2)	0.00181	(-0.6326, 3)	0.00321
(-0.5574, 1)	0.00777	(-0.5574, 2)	0.00163	(-0.5574, 3)	0.00462
(-0.5478, 1)	0.00752	(-0.5478, 2)	0.00171	(-0.5478, 3)	0.00303
(-0.5237, 1)	0.00287	(-0.5237, 2)	0.00172	(-0.5237, 3)	0.00281
(-0.4982, 1)	0.00255	(-0.4982, 2)	0.00148	(-0.4982, 3)	0.00196
(-0.4631, 1)	0.00195	(-0.4631, 2)	0.00166	(-0.4631, 3)	0.00400
(-0.4336, 1)	0.00170	(-0.4336, 2)	0.00187	(-0.4336, 3)	0.00272
(-0.3709, 1)	0.00164	(-0.3709, 2)	0.00197	(-0.3709, 3)	0.00193
(-0.2798, 1)	0.00154	(-0.2798, 2)	0.00218	(-0.2798, 3)	0.00259
(-0.1939, 1)	0.00159	(-0.1939, 2)	0.00298	(-0.1939, 3)	0.00199
(-0.1918, 1)	0.00143	(-0.1918, 2)	0.00326	(-0.1918, 3)	0.00289
(-0.1837, 1)	0.00205	(-0.1837, 2)	0.00556	(-0.1837, 3)	0.00211
(-0.1450, 1)	0.00173	(-0.1450, 2)	0.00760	(-0.1450, 3)	0.00170
(-0.1430, 1)	0.00161	(-0.1430, 2)	0.00568	(-0.1430, 3)	0.00128
(-0.1151, 1)	0.00423	(-0.1151, 2)	0.00220	(-0.1151, 3)	0.01389
(-0.0884, 1)	0.00351	(-0.0884, 2)	0.00175	(-0.0884, 3)	0.02066
(-0.0560, 1)	0.00514	(-0.0560, 2)	0.00176	(-0.0560, 3)	0.01154
(-0.0124, 1)	0.00380	(-0.0124, 2)	0.00214	(-0.0124, 3)	0.01050
(0.0000, 1)	0.00296	(0.0000, 2)	0.00284	(0.0000, 3)	0.00881
(0.0036, 1)	0.05584	(0.0036, 2)	0.00492	(0.0036, 3)	0.00569
(0.0350, 1)	0.06987	(0.0350, 2)	0.00677	(0.0350, 3)	0.00355
(0.0879, 1)	0.03947	(0.0879, 2)	0.00821	(0.0879, 3)	0.00258
(0.1222, 1)	0.03186	(0.1222, 2)	0.01139	(0.1222, 3)	0.00189
(0.2192, 1)	0.02311	(0.2192, 2)	0.01434	(0.2192, 3)	0.00215
(0.2223, 1)	0.01459	(0.2223, 2)	0.00523	(0.2223, 3)	0.00155
(0.2748, 1)	0.01459	(0.2748, 2)	0.00523	(0.2748, 3)	0.00155
(0.3154, 1)	0.01221	(0.3154, 2)	0.00334	(0.3154, 3)	0.00129
(0.4112, 1)	0.01151	(0.4112, 2)	0.00458	(0.4112, 3)	0.00129
(0.4434, 1)	0.01184	(0.4434, 2)	0.00738	(0.4434, 3)	0.00115
(0.4484, 1)	0.00883	(0.4484, 2)	0.01028	(0.4484, 3)	0.00105
(0.4677, 1)	0.00626	(0.4677, 2)	0.01178	(0.4677, 3)	0.00102
(0.5505, 1)	0.01574	(0.5505, 2)	0.01142	(0.5505, 3)	0.00159
(0.5662, 1)	0.02498	(0.5662, 2)	0.02024	(0.5662, 3)	0.00177
(0.7154, 1)	0.01499	(0.7154, 2)	0.03736	(0.7154, 3)	0.00173
(0.8502, 1)	0.02341	(0.8502, 2)	0.0229	(0.8502, 3)	0.00286
(1.1930, 1)	0.01985	(1.1930, 2)	0.00789	(1.1930, 3)	0.01769

Figure 1: Marginal Posterior Density of r




Table 2: Summary of Joint Posterior Densities of (d, r) for Chemical Data

(d, r)	Integrated Density	(d, r)	Integrated Density	(d, r)	Integrated Density
(1, -0.4)	0.03665	(2, -0.4)	0.01669	(2, -0.4)	0.00088
(1, -0.3)	0.15757	(2, -0.3)	0.01319	(2, -0.3)	0.00061
(1, -0.2)	0.15757	(2, -0.2)	0.01319	(2, -0.2)	0.00061
(1, -0.1)	0.09583	(2, -0.1)	0.00858	(2, -0.1)	0.00031
(1, 0.0)	0.17483	(2, 0.0)	0.01402	(2, 0.0)	0.00030
(1, 0.1)	0.04738	(2, 0.1)	0.02424	(2, 0.1)	0.00026
(1, 0.2)	0.03386	(2, 0.2)	0.02145	(2, 0.2)	0.00064
(1, 0.3)	0.03344	(2, 0.3)	0.01352	(2, 0.3)	0.00304
(1, 0.4)	0.09726	(2, 0.4)	0.00675	(2, 0.4)	0.02733

Smadi, M. M. (1997). *Bayesian inference of threshold autoregressive moving average models using sample-based methods*. Unpublished Ph. D. Dissertation, Colorado State University, Fort Collins, CO, U.S.A.

Tong H. (1990). *Nonlinear Time series models: A dynamical system approach*. London: Oxford University Press.

Tong, H., & Lim, K. S. (1980). Threshold autoregression, limit cycles and cyclical data. *Journal of the Royal Statistical Society, B42*, 245-92.

Wang, W. Y., Du, J. G., & Xiang, J. T. (1984). Threshold autoregressive moving average model. *Journal of Computational Mathematics, 4*, 414-19 (in Chinese).